

FGF19 antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP22412a

Product Information

Application	WB, IHC-P, E
Primary Accession	O95750
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit Ig
Clone Names	R03803
Calculated MW	24003

Additional Information

Gene ID	9965
Other Names	Fibroblast growth factor 19, FGF-19, FGF19
Target/Specificity	This antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between amino acids from human.
Dilution	WB~~1:1000 IHC-P~~1ug/ml E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	FGF19 antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	FGF19
Function	Involved in the suppression of bile acid biosynthesis through down-regulation of CYP7A1 expression, following positive regulation of the JNK and ERK1/2 cascades. Stimulates glucose uptake in adipocytes. Activity requires the presence of KLB and FGFR4.
Cellular Location	Secreted.
Tissue Location	Expressed in fetal brain, cartilage, retina, and adult gall bladder.


Background

Involved in the suppression of bile acid biosynthesis through down-regulation of CYP7A1 expression, following positive regulation of the JNK and ERK1/2 cascades. Stimulates glucose uptake in adipocytes. Activity requires the presence of KLB and FGFR4.


References

Nishimura T.,et al.Biochim. Biophys. Acta 1444:148-151(1999).
Xie M.-H.,et al.Cytokine 11:729-735(1999).
Clark H.F.,et al.Genome Res. 13:2265-2270(2003).
Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.
Zhang Z.,et al.Protein Sci. 13:2819-2824(2004).

Images



All lanes: Anti-FGF19 antibody at 1:1000 dilution + Mouse liver lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 24 KDa Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemical analysis of paraffin-embedded Human liver cancer tissue using FGF19 antibody performed on the Abcarta® FAIP-48T Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9. 0). Samples were incubated with primary antibody(Ready-to-use) for 15 min at room temperature. AmpSee™ Detection Systems(Abcepta:ADR005) was used as the secondary antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.