

Gasdermin-E antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP22465a

Product Information

Application	WB, E
Primary Accession	O60443
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit Ig
Clone Names	R03173
Calculated MW	54555

Additional Information

Gene ID	1687
Other Names	Gasdermin-E, Inversely correlated with estrogen receptor expression 1, ICERE-1, Non-syndromic hearing impairment protein 5, Gasdermin-E, N-terminal, GSDME-NT, Gasdermin-E, C-terminal, GSDME-CT, GSDME {ECO:0000303 PubMed:28459430, ECO:0000312 HGNC:HGNC:2810}
Target/Specificity	This antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between amino acids from human.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Gasdermin-E antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	GSDME {ECO:0000303 PubMed:28459430, ECO:0000312 HGNC:HGNC:2810}
Function	[Gasdermin-E]: Precursor of a pore-forming protein that converts non-inflammatory apoptosis to pyroptosis (PubMed: 27281216 , PubMed: 28459430 , PubMed: 33852854 , PubMed: 35594856 , PubMed: 36607699). This form constitutes the precursor of the pore-forming

protein: upon cleavage, the released N-terminal moiety (Gasdermin-E, N-terminal) binds to membranes and forms pores, triggering pyroptosis (PubMed:[28459430](#)).

Cellular Location	[Gasdermin-E, N-terminal]: Cell membrane; Multi-pass membrane protein {ECO:0000250 UniProtKB:Q5Y4Y6}
Tissue Location	Expressed in cochlea (PubMed:9771715). Low level of expression in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas, with highest expression in placenta (PubMed:9771715).


Background

[Gasdermin-E]: Precursor of a pore-forming protein that converts non-inflammatory apoptosis to pyroptosis (PubMed:[27281216](#), PubMed:[28459430](#), PubMed:[33852854](#), PubMed:[35594856](#), PubMed:[36607699](#)). This form constitutes the precursor of the pore-forming protein: upon cleavage, the released N-terminal moiety (Gasdermin-E, N-terminal) binds to membranes and forms pores, triggering pyroptosis (PubMed:[28459430](#)).

References

Van Laer L.,et al.Nat. Genet. 20:194-197(1998).
Van Laer L.,et al.Submitted (JUN-1998) to the EMBL/GenBank/DDBJ databases.
Mei G.,et al.Submitted (FEB-1999) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Hillier L.W.,et al.Nature 424:157-164(2003).

Images



All lanes: Anti-Gasdermin-E antibody at 1:1000 dilution
Lane 1: Rat brain lysate Lane 2: U-87 MG whole cell lysate
Lane 3: MCF-7 whole cell lysate Lane 4: U-251 MG whole cell lysate
Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 55 KDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.