

ACTA2 antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP22466a

Product Information

Application	WB, E
Primary Accession	P62736
Reactivity	Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit Ig
Clone Names	R03324
Calculated MW	42009

Additional Information

Gene ID	59
Other Names	Actin, aortic smooth muscle, 3.6.4.-, Alpha-actin-2, Cell growth-inhibiting gene 46 protein, Actin, aortic smooth muscle, intermediate form, ACTA2, ACTSA, ACTVS
Target/Specificity	This antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between amino acids from human.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ACTA2 antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ACTA2
Synonyms	ACTSA, ACTVS
Function	Actins are highly conserved proteins that are involved in various types of cell motility and are ubiquitously expressed in all eukaryotic cells.

Background

Actins are highly conserved proteins that are involved in various types of cell motility and are ubiquitously expressed in all eukaryotic cells.

References

Kamada S.,et al.Nucleic Acids Res. 17:1767-1767(1989).


Reddy S.,et al.J. Biol. Chem. 265:1683-1687(1990).

Kim J.W.,et al.Submitted (JUL-2004) to the EMBL/GenBank/DBJ databases.

Halleck A.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DBJ databases.

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Images



All lanes: Anti-ACTA2 antibody at 1:2000 dilution Lane 1: C2C12 whole cell lysate Lane 2: NIH/3T3 whole cell lysate Lane 3: Mouse uterus lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 42 KDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.