

hIL1R-pY496

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP22476a

Product Information

Application	WB, E
Primary Accession	P14778
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit Ig
Clone Names	R04087NP
Calculated MW	65402

Additional Information

Gene ID	3554
Other Names	Interleukin-1 receptor type 1, IL-1R-1, IL-1RT-1, IL-1RT1, 3.2.2.6 {ECO:0000255 PROSITE-ProRule:PRU00204}, CD121 antigen-like family member A, Interleukin-1 receptor alpha, IL-1R-alpha, Interleukin-1 receptor type I, p80, CD121a, Interleukin-1 receptor type 1, membrane form, mIL-1R1, mIL-1RI, Interleukin-1 receptor type 1, soluble form, sIL-1R1, sIL-1RI, IL1R1, IL1R, IL1RA, IL1RT1
Target/Specificity	This antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between amino acids from human.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	hIL1R-pY496 is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	IL1R1
Synonyms	IL1R, IL1RA, IL1RT1

Function	Receptor for IL1A, IL1B and IL1RN (PubMed: 2950091 , PubMed: 37315560). After binding to interleukin-1 associates with the coreceptor IL1RAP to form the high affinity interleukin-1 receptor complex which mediates interleukin-1-dependent activation of NF-kappa- B, MAPK and other pathways. Signaling involves the recruitment of adapter molecules such as TOLLIP, MYD88, and IRAK1 or IRAK2 via the respective TIR domains of the receptor/coreceptor subunits. Binds ligands with comparable affinity and binding of antagonist IL1RN prevents association with IL1RAP to form a signaling complex. Involved in IL1B-mediated costimulation of IFNG production from T-helper 1 (Th1) cells (PubMed: 10653850).
Cellular Location	Membrane; Single- pass type I membrane protein. Cell membrane. Secreted
Tissue Location	Expressed in T-helper cell subsets. Preferentially expressed in T-helper 1 (Th1) cells.


Background

Receptor for IL1A, IL1B and IL1RN (PubMed: [2950091](#), PubMed:[37315560](#)). After binding to interleukin-1 associates with the coreceptor IL1RAP to form the high affinity interleukin-1 receptor complex which mediates interleukin-1-dependent activation of NF-kappa- B, MAPK and other pathways. Signaling involves the recruitment of adapter molecules such as TOLLIP, MYD88, and IRAK1 or IRAK2 via the respective TIR domains of the receptor/coreceptor subunits. Binds ligands with comparable affinity and binding of antagonist IL1RN prevents association with IL1RAP to form a signaling complex. Involved in IL1B-mediated costimulation of IFNG production from T-helper 1 (Th1) cells (PubMed:[10653850](#)).

References

Chua A.O.,et al.Nucleic Acids Res. 17:10114-10114(1989).
Sims J.E.,et al.Proc. Natl. Acad. Sci. U.S.A. 86:8946-8950(1989).
Hillier L.W.,et al.Nature 434:724-731(2005).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Mosley B.,et al.J. Biol. Chem. 262:2941-2944(1987).

Images



All lanes: Anti-IL1R Antibody pY496 at 1:2000 dilution +
Mouse lung lysate Lysates/proteins at 20 µg per lane.
Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase
conjugated (ASP1615) at 1/15000 dilution. Observed band
size: 65 KDa Blocking/Dilution buffer: 5% NFDm/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.