

Endophilin-pY80

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP22486a

Product Information

Application	WB, E
Primary Accession	Q9Y371
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit Ig
Clone Names	R05012NP
Calculated MW	40796

Additional Information

Gene ID	51100
Other Names	Endophilin-B1, Bax-interacting factor 1, Bif-1, SH3 domain-containing GRB2-like protein B1, SH3GLB1, KIAA0491
Target/Specificity	This antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between amino acids from human.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Endophilin-pY80 is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SH3GLB1
Synonyms	KIAA0491
Function	May be required for normal outer mitochondrial membrane dynamics (PubMed: 15452144). Required for coatomer-mediated retrograde transport in certain cells (By similarity). May recruit other proteins to membranes with high curvature. May promote membrane fusion (PubMed: 11604418). Involved

in activation of caspase-dependent apoptosis by promoting BAX/BAK1 activation (PubMed:[16227588](#)). Isoform 1 acts proapoptotic in fibroblasts (By similarity). Involved in caspase- independent apoptosis during nutrition starvation and involved in the regulation of autophagy. Activates lipid kinase activity of PIK3C3 during autophagy probably by associating with the PI3K complex II (PI3KC3-C2) (PubMed:[17891140](#)). Associated with PI3KC3-C2 during autophagy may regulate the trafficking of ATG9A from the Golgi complex to the peripheral cytoplasm for the formation of autophagosomes by inducing Golgi membrane tubulation and fragmentation (PubMed:[21068542](#)). Involved in regulation of degradative endocytic trafficking and cytokinesis, probably in the context of PI3KC3-C2 (PubMed:[20643123](#)). Isoform 2 acts antiapoptotic in neuronal cells; involved in maintenance of mitochondrial morphology and promotes neuronal viability (By similarity).

Cellular Location

Cytoplasm. Golgi apparatus membrane; Peripheral membrane protein. Mitochondrion outer membrane; Peripheral membrane protein. Cytoplasmic vesicle, autophagosome membrane. Midbody. Note=Association with the Golgi apparatus depends on the cell type (By similarity). Following starvation colocalizes with ATG5 and LC3 autophagy-related protein(s) on autophagosomal membranes (PubMed:17891140). {ECO:0000250, ECO:0000269 | PubMed:17891140}

Tissue Location

Highly expressed in heart, skeletal muscle, kidney and placenta. Detected at lower levels in brain, colon, thymus, spleen, liver, small intestine, lung and peripheral blood leukocytes

Background


May be required for normal outer mitochondrial membrane dynamics (PubMed: [15452144](#)). Required for coatamer-mediated retrograde transport in certain cells (By similarity). May recruit other proteins to membranes with high curvature. May promote membrane fusion (PubMed:[11604418](#)). Involved in activation of caspase-dependent apoptosis by promoting BAX/BAK1 activation (PubMed:[16227588](#)). Isoform 1 acts proapoptotic in fibroblasts (By similarity). Involved in caspase- independent apoptosis during nutrition starvation and involved in the regulation of autophagy. Activates lipid kinase activity of PIK3C3 during autophagy probably by associating with the PI3K complex II (PI3KC3-C2) (PubMed:[17891140](#)). Associated with PI3KC3-C2 during autophagy may regulate the trafficking of ATG9A from the Golgi complex to the peripheral cytoplasm for the formation of autophagosomes by inducing Golgi membrane tubulation and fragmentation (PubMed:[21068542](#)). Involved in regulation of degradative endocytic trafficking and cytokinesis, probably in the context of PI3KC3-C2 (PubMed:[20643123](#)). Isoform 2 acts antiapoptotic in neuronal cells; involved in maintenance of mitochondrial morphology and promotes neuronal viability (By similarity).

References

Pierrat B.,et al.Genomics 71:222-234(2001).
 Cuddeback S.M.,et al.J. Biol. Chem. 276:20559-20565(2001).
 Modregger J.,et al.J. Biol. Chem. 278:4160-4167(2003).
 Seki N.,et al.DNA Res. 4:345-349(1997).
 Lai C.-H.,et al.Genome Res. 10:703-713(2000).

Images

All lanes: Anti-Endophilin-pY80 at 1:1000 dilution + K562 whole cell lysate Lysates/proteins at 20 µg per lane.
 Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase



conjugated (ASP1615) at 1/15000 dilution. Observed band size: 41 KDa Blocking/Dilution buffer: 5% NFDm/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.