

SHP2(Y584) Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP22489a

Product Information

Application	WB, E
Primary Accession	Q06124
Other Accession	P35235 , P41499
Reactivity	Human
Predicted	Mouse, Anthrax LF (Lethal Factor) produced by Bacillus an
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit Ig
Clone Names	R05330NP
Calculated MW	68011

Additional Information

Gene ID	5781
Other Names	Tyrosine-protein phosphatase non-receptor type 11, 3.1.3.48, Protein-tyrosine phosphatase 1D, PTP-1D, Protein-tyrosine phosphatase 2C, PTP-2C, SH-PTP2, SHP-2, Shp2, SH-PTP3, PTPN11, PTP2C, SHPTP2
Target/Specificity	This SHP2(Y584) antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between amino acids from the human region of human SHP2(Y584).
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	SHP2(Y584) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PTPN11
Synonyms	PTP2C, SHPTP2

Function	Acts downstream of various receptor and cytoplasmic protein tyrosine kinases to participate in the signal transduction from the cell surface to the nucleus (PubMed: 10655584 , PubMed: 14739280 , PubMed: 18559669 , PubMed: 18829466 , PubMed: 26742426 , PubMed: 28074573 , PubMed: 32184441). Positively regulates MAPK signal transduction pathway (PubMed: 28074573). Dephosphorylates GAB1, ARHGAP35 and EGFR (PubMed: 28074573). Dephosphorylates ROCK2 at 'Tyr-722' resulting in stimulation of its RhoA binding activity (PubMed: 18559669). Dephosphorylates CDC73 (PubMed: 26742426). Dephosphorylates SOX9 on tyrosine residues, leading to inactivate SOX9 and promote ossification (By similarity). Dephosphorylates tyrosine-phosphorylated NEDD9/CAS-L (PubMed: 19275884). Acts as an effector of PDCD1-mediated inhibition of T-cell response: recruited by phosphorylated PDCD1, mediating dephosphorylation of key T-cell receptor (TCR) proximal signaling molecules, leading to TCR signaling inhibition (PubMed: 32184441).
Cellular Location	Cytoplasm. Nucleus
Tissue Location	Widely expressed, with highest levels in heart, brain, and skeletal muscle.


Background

Acts downstream of various receptor and cytoplasmic protein tyrosine kinases to participate in the signal transduction from the cell surface to the nucleus (PubMed:[10655584](#), PubMed:[14739280](#), PubMed:[18559669](#), PubMed:[18829466](#), PubMed:[26742426](#), PubMed:[28074573](#), PubMed:[32184441](#)). Positively regulates MAPK signal transduction pathway (PubMed:[28074573](#)). Dephosphorylates GAB1, ARHGAP35 and EGFR (PubMed:[28074573](#)). Dephosphorylates ROCK2 at 'Tyr-722' resulting in stimulation of its RhoA binding activity (PubMed:[18559669](#)). Dephosphorylates CDC73 (PubMed:[26742426](#)). Dephosphorylates SOX9 on tyrosine residues, leading to inactivate SOX9 and promote ossification (By similarity). Dephosphorylates tyrosine-phosphorylated NEDD9/CAS-L (PubMed:[19275884](#)). Acts as an effector of PDCD1-mediated inhibition of T-cell response: recruited by phosphorylated PDCD1, mediating dephosphorylation of key T-cell receptor (TCR) proximal signaling molecules, leading to TCR signaling inhibition (PubMed:[32184441](#)).

References

Adachi M.,et al.FEBS Lett. 314:335-339(1992).
Freeman R.M. Jr.,et al.Proc. Natl. Acad. Sci. U.S.A. 89:11239-11243(1992).
Bastien L.,et al.Biochem. Biophys. Res. Commun. 196:124-133(1993).
Ahmad S.,et al.Proc. Natl. Acad. Sci. U.S.A. 90:2197-2201(1993).
Vogel W.,et al.Science 259:1611-1614(1993).

Images



All lanes: Anti-SHP2(Y584) Antibody at 1:2000 dilution + A431 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 68 KDa Blocking/Dilution buffer: 5% NFDm/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.