

E2F1(S332) Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP22491a

Product Information

Application	WB, E
Primary Accession	Q01094
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit Ig
Clone Names	R05606NP
Calculated MW	46920

Additional Information

Gene ID	1869
Other Names	Transcription factor E2F1, E2F-1, PBR3, Retinoblastoma-associated protein 1, RBAP-1, Retinoblastoma-binding protein 3, RBBP-3, pRB-binding protein E2F-1, E2F1 {ECO:0000303 PubMed:8964493, ECO:0000312 HGNC:HGNC:3113}
Target/Specificity	This E2F1(S332) antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between amino acids from the human region of human E2F1(S332).
Dilution	WB~~1:500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	E2F1(S332) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	E2F1 {ECO:0000303 PubMed:8964493, ECO:0000312 HGNC:HGNC:3113}
Function	Transcription activator that binds DNA cooperatively with DP proteins through the E2 recognition site, 5'-TTTC[CG]CGC-3' found in the promoter region of a number of genes whose products are involved in cell cycle

regulation or in DNA replication (PubMed:[10675335](#), PubMed:[12717439](#), PubMed:[17050006](#), PubMed:[17704056](#), PubMed:[18625225](#), PubMed:[28992046](#)). The DRTF1/E2F complex functions in the control of cell-cycle progression from G1 to S phase (PubMed:[10675335](#), PubMed:[12717439](#), PubMed:[17704056](#)). E2F1 binds preferentially RB1 in a cell-cycle dependent manner (PubMed:[10675335](#), PubMed:[12717439](#), PubMed:[17704056](#)). It can mediate both cell proliferation and TP53/p53-dependent apoptosis (PubMed:[8170954](#)). Blocks adipocyte differentiation by binding to specific promoters repressing CEBPA binding to its target gene promoters (PubMed:[20176812](#)). Directly activates transcription of PEG10 (PubMed:[17050006](#), PubMed:[18625225](#), PubMed:[28992046](#)). Positively regulates transcription of RRP1B (PubMed:[20040599](#)).

Cellular Location Nucleus


Background

Transcription activator that binds DNA cooperatively with DP proteins through the E2 recognition site, 5'-TTTC[CG]CGC-3' found in the promoter region of a number of genes whose products are involved in cell cycle regulation or in DNA replication (PubMed:[10675335](#), PubMed:[12717439](#), PubMed:[17050006](#), PubMed:[17704056](#), PubMed:[18625225](#), PubMed:[28992046](#)). The DRTF1/E2F complex functions in the control of cell-cycle progression from G1 to S phase (PubMed:[10675335](#), PubMed:[12717439](#), PubMed:[17704056](#)). E2F1 binds preferentially RB1 in a cell-cycle dependent manner (PubMed:[10675335](#), PubMed:[12717439](#), PubMed:[17704056](#)). It can mediate both cell proliferation and TP53/p53- dependent apoptosis (PubMed:[8170954](#)). Blocks adipocyte differentiation by binding to specific promoters repressing CEBPA binding to its target gene promoters (PubMed:[20176812](#)). Directly activates transcription of PEG10 (PubMed:[17050006](#), PubMed:[18625225](#), PubMed:[28992046](#)). Positively regulates transcription of RRP1B (PubMed:[20040599](#)).

References

Helin K.,et al.Cell 70:337-350(1992).
Kaelin W.G. Jr.,et al.Cell 70:351-364(1992).
Shan B.,et al.Mol. Cell. Biol. 12:5620-5631(1992).
Neuman E.,et al.Gene 173:163-169(1996).
Deloukas P.,et al.Nature 414:865-871(2001).

Images



All lanes: Anti-E2F1(S332) Antibody at 1:500 dilution Lane 1: SH-SY5Y whole cell lysate Lane 2: C6 whole cell lysate Lane 3: U-2OS whole cell lysate Lane 4: MCF-7 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 60 KDa Blocking/Dilution buffer: 5% NFDm/TBST.