

RNF36 (TRIM69) Antibody (Center H215)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP2525c

Product Information

Application	WB, E
Primary Accession	Q86WT6
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB13950
Calculated MW	57419
Antigen Region	200-231

Additional Information

Gene ID	140691
Other Names	E3 ubiquitin-protein ligase TRIM69, 632-, RFP-like domain-containing protein trimless, RING finger protein 36, Tripartite motif-containing protein 69, TRIM69, RNF36
Target/Specificity	This RNF36 (TRIM69) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 200-231 amino acids from the Central region of human RNF36 (TRIM69).
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	RNF36 (TRIM69) Antibody (Center H215) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TRIM69
Synonyms	RNF36
Function	E3 ubiquitin ligase that plays an important role in antiviral immunity by

restricting different viral infections including dengue virus or vesicular stomatitis indiana virus (PubMed:[23131556](#), PubMed:[30142214](#), PubMed:[31375575](#), PubMed:[31578292](#)). Ubiquitinates viral proteins such as dengue virus NS3 thereby limiting infection (PubMed:[30844644](#)). In addition, acts as a key mediator of type I interferon induced microtubule stabilization by directly associating to microtubules independently of its E3 ligase activity (PubMed:[36251989](#)). Also plays a role in cataract formation together with TP53 (PubMed:[30844644](#)). Mechanistically, inhibits UVB-induced cell apoptosis and reactive oxygen species (ROS) production by inducing TP53 ubiquitination (PubMed:[30844644](#)). Regulates centrosome dynamics and mitotic progression by ubiquitinating STK3/MST2; leading to its redistribution to the perinuclear cytoskeleton and subsequent phosphorylation by PLK1 (PubMed:[37739411](#)).

Cellular Location

Cytoplasm. Nucleus. Nucleus speckle. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Note=Adopts a filamentous distribution in the cell cytoplasm where it strongly colocalizes with stable microtubules


Background

TRIM69 is a member of the RING-B-box-coiled-coil (RBCC) family with an N-terminal RING finger motif, a PRY domain and a C-terminal SPRY domain. The mouse ortholog of TRIM69 is specifically expressed in germ cells at the round spermatid stages during spermatogenesis and, when overexpressed, induces apoptosis.


References

Shyu,H.W., Exp. Cell Res. 287 (2), 301-313 (2003)
Shyu,H.W., Mech. Dev. 108 (1-2), 213-216 (2001)

Images



Anti-TRIM69 Antibody (Center H215) at 1:1000 dilution + PC-3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 57 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot analysis of anti-TRIM69 Antibody (Center H215) (RB13950) in K562 cell line lysates (35µg/lane). TRIM69(arrow) was detected using the purified Pab.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.