

RNF36 (TRIM69) Antibody (Center H215)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP2525c

Product Information

Application WB, E **Primary Accession Q86WT6** Reactivity Human Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Clone Names** RB13950 **Calculated MW** 57419 **Antigen Region** 200-231

Additional Information

Gene ID 140691

Other Names E3 ubiquitin-protein ligase TRIM69, 632-, RFP-like domain-containing protein

trimless, RING finger protein 36, Tripartite motif-containing protein 69,

TRIM69, RNF36

Target/Specificity This RNF36 (TRIM69) antibody is generated from rabbits immunized with a

KLH conjugated synthetic peptide between 200-231 amino acids from the

Central region of human RNF36 (TRIM69).

Dilution WB~~1:1000 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation

followed by dialysis against PBS.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions RNF36 (TRIM69) Antibody (Center H215) is for research use only and not for

use in diagnostic or therapeutic procedures.

Protein Information

Name TRIM69

Synonyms RNF36

Function E3 ubiquitin ligase that plays an important role in antiviral immunity by

restricting different viral infections including dengue virus or vesicular stomatitis indiana virus (PubMed:23131556, PubMed:30142214, PubMed:31375575, PubMed:31578292). Ubiquitinates viral proteins such as dengue virus NS3 thereby limiting infection (PubMed:30844644). In addition, acts as a key mediator of type I interferon induced microtubule stabilization by directly associating to microtubules independently of its E3 ligase activity (PubMed:36251989). Also plays a role in cataract formation together with TP53 (PubMed:30844644). Mechanistically, inhibits UVB-induced cell apoptosis and reactive oxygen species (ROS) production by inducing TP53 ubiquitination (PubMed:30844644). Regulates centrosome dynamics and mitotic progression by ubiquitinating STK3/MST2; leading to its redistribution to the perinuclear cytoskeleton and subsequent phosphorylation by PLK1 (PubMed:37739411).

Cellular Location

Cytoplasm. Nucleus. Nucleus speckle. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Note=Adopts a filamentous distribution in the cell cytoplasm where it strongly colocalizes with stable microtubules


Background

TRIM69 is a member of the RING-B-box-coiled-coil (RBCC) family with an N-terminal RING finger motif, a PRY domain and a C-terminal SPRY domain. The mouse ortholog of TRIM69 is specifically expressed in germ cells at the round spermatid stages during spermatogenesis and, when overexpressed, induces apoptosis.


References

Shyu,H.W., Exp. Cell Res. 287 (2), 301-313 (2003) Shyu,H.W., Mech. Dev. 108 (1-2), 213-216 (2001)

Images



Anti-TRIM69 Antibody (Center H215) at 1:1000 dilution + PC-3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 57 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot analysis of anti-TRIM69 Antibody (Center H215) (RB13950) in K562 cell line lysates (35ug/lane). TRIM69(arrow) was detected using the purified Pab.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.