

# DAX1 (NR0B1) Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP2708a

## Product Information

---

Application	WB, E
Primary Accession	<a href="#">P51843</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	51718
Antigen Region	4-34

## Additional Information

---

Gene ID	190
Other Names	Nuclear receptor subfamily 0 group B member 1, DSS-AHC critical region on the X chromosome protein 1, Nuclear receptor DAX-1, NR0B1, AHC, DAX1
Target/Specificity	This DAX1 (NR0B1) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 4-34 amino acids from the N-terminal region of human DAX1 (NR0B1).
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	DAX1 (NR0B1) Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	NR0B1
Synonyms	AHC, DAX1 {ECO:0000303 PubMed:26416531}
Function	Nuclear receptor that lacks a DNA-binding domain and acts as a corepressor that inhibits the transcriptional activity of other nuclear receptors through heterodimeric interactions (PubMed: <a href="#">12482977</a> , PubMed: <a href="#">32433991</a> ).

Component of a cascade required for the development of the hypothalamic-pituitary-adrenal-gonadal axis (PubMed:[7990953](#), PubMed:[8675564](#)). May also have a role in the development of the embryo and in the maintenance of embryonic stem cell pluripotency (By similarity).

#### Cellular Location

Nucleus. Cytoplasm. Note=Shuttles between the cytoplasm and nucleus. Homodimers exits in the cytoplasm and in the nucleus


## Background

NROB1 contains a DNA-binding domain and acts as a dominant-negative regulator of transcription which is mediated by the retinoic acid receptor. This protein also functions as an anti-testis gene by acting antagonistically to Sry. Mutations in the gene for NROB1 result in both X-linked congenital adrenal hypoplasia and hypogonadotropic hypogonadism.


## References

Calliari,L.E., Genet. Mol. Res. 6 (2), 177-183 (2007)  
Kinsey,M., Mol. Cancer Res. 4 (11), 851-859 (2006)

## Images



All lanes : Anti-NROB1 Antibody (N-term) at 1:1000 dilution Lane 1: HeLa whole cell lysate Lane 2: NCI-H460 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 52 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



NROB1 Antibody (N-term) (Cat. #AP2708a) western blot analysis in HeLa,NCI-H292,PC-3 cell line lysates (35ug/lane).This demonstrates the NROB1 antibody detected the NROB1 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.