10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

PHB Antibody (C-term E194)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2710d

Product Information

Application	WB, IHC-P, FC, E
Primary Accession	P35232
Other Accession	P67779, P67778, Q3T165
Reactivity	Human
Predicted	Bovine, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB17564
Calculated MW	29804
Antigen Region	179-205

Additional Information

Gene ID	5245
Other Names	Prohibitin, PHB
Target/Specificity	This PHB antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 179-205 amino acids from the C-terminal region of human PHB.
Dilution	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PHB Antibody (C-term E194) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PHB1 {ECO:0000303|PubMed:28017329, ECO:0000312|HGNC:HGNC:8912}
Function	Protein with pleiotropic attributes mediated in a cell- compartment- and tissue-specific manner, which include the plasma membrane-associated cell

signaling functions, mitochondrial chaperone, and transcriptional co-regulator of transcription factors in the nucleus (PubMed:11302691, PubMed:20959514, PubMed:28017329, PubMed:31522117). Plays a role in adipose tissue and glucose homeostasis in a sex-specific manner (By similarity). Contributes to pulmonary vascular remodeling by accelerating proliferation of pulmonary arterial smooth muscle cells (By similarity).

Cellular Location Mitochondrion inner membrane. Nucleus. Cytoplasm. Cell membrane

Tissue Location Widely expressed in different tissues.


Background

Prohibitin is an evolutionarily conserved protein that is ubiquitously expressed. It is thought to be a negative regulator of cell proliferation and may be a tumor suppressor. Mutations have been linked to sporadic breast cancer. Prohibitin is expressed as two transcripts with varying lengths of 3' untranslated region.


References

Gregory-Bass,R.C., Int. J. Cancer 122 (9), 1923-1930 (2008)
Ross,J.A., J. Biol. Chem. 283 (8), 4699-4713 (2008)
White,J.J., Genomics 11 (1), 228-230 (1991)

Images




Western blot analysis of PHB (arrow) using rabbit polyclonal PHB Antibody (C-term) (Cat.#AP2710d). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the PHB gene (Lane 2) (Origene Technologies).




Western blot analysis of anti-PHB Antibody (C-term) (Cat.#AP2710d) in Jurkat cell line lysates (35ug/lane).PHB (arrow) was detected using the purified Pab.

Formalin-fixed and paraffin-embedded human breast carcinoma reacted with PHB Antibody (C-term E194), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been



evaluated.



Flow cytometric analysis of NCI-H292 cells using PHB Antibody (C-term E194)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.