

LGR5 (GPR49) Antibody (loop1)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP2745c

Product Information

Application	IHC-P, WB, E
Primary Accession	O75473
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB14210
Calculated MW	99998
Antigen Region	605-638

Additional Information

Gene ID	8549
Other Names	Leucine-rich repeat-containing G-protein coupled receptor 5, G-protein coupled receptor 49, G-protein coupled receptor 67, G-protein coupled receptor HG38, LGR5, GPR49, GPR67
Target/Specificity	This LGR5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 605-638 amino acids from human LGR5.
Dilution	IHC-P~~1:100~500 WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	LGR5 (GPR49) Antibody (loop1) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	LGR5
Synonyms	GPR49, GPR67
Function	Receptor for R-spondins that potentiates the canonical Wnt signaling

pathway and acts as a stem cell marker of the intestinal epithelium and the hair follicle. Upon binding to R-spondins (RSPO1, RSPO2, RSPO3 or RSPO4), associates with phosphorylated LRP6 and frizzled receptors that are activated by extracellular Wnt receptors, triggering the canonical Wnt signaling pathway to increase expression of target genes. In contrast to classical G-protein coupled receptors, does not activate heterotrimeric G-proteins to transduce the signal. Involved in the development and/or maintenance of the adult intestinal stem cells during postembryonic development.

Cellular Location

Cell membrane; Multi-pass membrane protein. Golgi apparatus, trans-Golgi network membrane; Multi-pass membrane protein Note=Rapidly and constitutively internalized to the trans-Golgi network at steady state. Internalization to the trans-Golgi network may be the result of phosphorylation at Ser-861 and Ser-864; however, the phosphorylation event has not been proven (PubMed:23439653)

Tissue Location

Expressed in skeletal muscle, placenta, spinal cord, and various region of brain. Expressed at the base of crypts in colonic and small mucosa stem cells. In premalignant cancer expression is not restricted to the crypt base. Overexpressed in cancers of the ovary, colon and liver.


Background

LGR5/GPR49 is an orphan receptor. It may be an important receptor for signals controlling growth and differentiation of specific embryonic tissues.

References


Barker,N., Nature 449 (7165), 1003-1007 (2007) McClanahan,T., Cancer Biol. Ther. 5 (4), 419-426 (2006) Yamamoto,Y., Hepatology 37 (3), 528-533 (2003) Hsu,S.Y., Mol. Endocrinol. 14 (8), 1257-1271 (2000)

Images



Formalin-fixed and paraffin-embedded human brain tissue reacted with LGR5/GPR49 antibody (loop1) (Cat.#AP2745c), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

LGR5/GPR49 Antibody (loop1) (Cat. #AP11867a) western blot analysis in HL-60 cell line lysates (35ug/lane).This demonstrates the LGR5/GPR49 antibody detected the LGR5/GPR49 protein (arrow).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.