10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

SSB Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2801b

Product Information

Application	WB, IHC-P, E
Primary Accession	P05455
Other Accession	Q04504
Reactivity	Human
Predicted	Rabbit, Canine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB17352
Calculated MW	46837
Antigen Region	366-397

Additional Information

Gene ID	6741
Other Names	Lupus La protein, La autoantigen, La ribonucleoprotein, Sjoegren syndrome type B antigen, SS-B, SSB
Target/Specificity	This SSB antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 366-397 amino acids from the C-terminal region of human SSB.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	SSB Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SSB
Function	Binds to the 3' poly(U) terminus of nascent RNA polymerase III transcripts, protecting them from exonuclease digestion and facilitating their folding and

maturation (PubMed:2470590, PubMed:3192525). In case of Coxsackievirus B3 infection, binds to the viral internal ribosome entry site (IRES) and stimulates the IRES- mediated translation (PubMed:12384597).

Cellular Location Nucleus.


Background

SSB is involved in diverse aspects of RNA metabolism, including binding and protecting 3-prime UUU(OH) elements of newly RNA polymerase III -transcribed RNA, processing 5-prime and 3-prime ends of pre-tRNA precursors, acting as an RNA chaperone, and binding viral RNAs associated with hepatitis C virus. This protein was originally defined by its reactivity with autoantibodies from patients with Sjogren syndrome and systemic lupus erythematosus.


References

Bitko,V., J. Virol. 82 (16), 7977-7987 (2008)
Kotik-Kogan,O., Structure 16 (6), 852-862 (2008)
Biswas,D., Clin. Rheumatol. 27 (6), 717-722 (2008)

Images



Western blot analysis of SSB Antibody (C-term) (Cat.#AP2801b) in K562, Hela, 293 cell line lysates (35ug/lane). SSB (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human hepatocarcinoma reacted with ELAVL1 Antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.