

# PSMD11 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP2803b

## Product Information

---

<b>Application</b>	WB, IHC-P, E
<b>Primary Accession</b>	<a href="#">O00231</a>
<b>Other Accession</b>	<a href="#">F1LMZ8</a> , <a href="#">Q8BG32</a> , <a href="#">Q2KI42</a>
<b>Reactivity</b>	Human, Mouse
<b>Predicted</b>	Bovine, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB17375
<b>Calculated MW</b>	47464
<b>Antigen Region</b>	288-317

## Additional Information

---

<b>Gene ID</b>	5717
<b>Other Names</b>	26S proteasome non-ATPase regulatory subunit 11, 26S proteasome regulatory subunit RPN6, 26S proteasome regulatory subunit S9, 26S proteasome regulatory subunit p445, PSMD11
<b>Target/Specificity</b>	This PSMD11 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 288-317 amino acids from the C-terminal region of human PSMD11.
<b>Dilution</b>	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	PSMD11 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	PSMD11
<b>Function</b>	Component of the 26S proteasome, a multiprotein complex involved in the

ATP-dependent degradation of ubiquitinated proteins. This complex plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins, which could impair cellular functions, and by removing proteins whose functions are no longer required. Therefore, the proteasome participates in numerous cellular processes, including cell cycle progression, apoptosis, or DNA damage repair. In the complex, PSMD11 is required for proteasome assembly. Plays a key role in increased proteasome activity in embryonic stem cells (ESCs): its high expression in ESCs promotes enhanced assembly of the 26S proteasome, followed by higher proteasome activity.

**Cellular Location**

Nucleus. Cytoplasm, cytosol

**Tissue Location**

Highly expressed in embryonic stem cells (ESCs). Expression decreases as ESCs differentiate

## Background

---

The 26S proteasome (PSMD11) is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. The protein is a non-ATPase subunit of the 19S regulator.


## References

---

Saito,A., Gene 203 (2), 241-250 (1997)  
Hoffman,L., FEBS Lett. 404 (2-3), 179-184 (1997)



## Images

---



Western blot analysis of PSMD11 Antibody (C-term) (Cat.#AP2803b) in 293, Jurkat cell line lysates (35ug/lane). PSMD11 (arrow) was detected using the purified Pab.

Formalin-fixed and paraffin-embedded human lung carcinoma tissue reacted with PSMD11 antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



PSMD11 Antibody (C-term) (Cat.#AP2803b) western blot analysis in mouse kidney tissue lysates (35ug/lane). This demonstrates the PSMD11 antibody detected the PSMD11 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.