

# ENPP2 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP2854D

## Product Information

---

Application	FC, WB, E
Primary Accession	<a href="#">Q13822</a>
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB18117
Calculated MW	98994
Antigen Region	377-406

## Additional Information

---

Gene ID	5168
Other Names	Ectonucleotide pyrophosphatase/phosphodiesterase family member 2, E-NPP 2, Autotaxin, Extracellular lysophospholipase D, LysoPLD, ENPP2, ATX, PDNP2
Target/Specificity	This ENPP2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 377-406 amino acids from the Central region of human ENPP2.
Dilution	FC~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ENPP2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	ENPP2 ( <a href="#">HGNC:3357</a> )
Function	Secreted lysophospholipase D that hydrolyzes lysophospholipids to produce the signaling molecule lysophosphatidic acid (LPA) in extracellular fluids (PubMed: <a href="#">12354767</a> , PubMed: <a href="#">14500380</a> , PubMed: <a href="#">15769751</a> , PubMed: <a href="#">26371182</a> , PubMed: <a href="#">27754931</a> ). Its major substrate is

lysophosphatidylcholine (PubMed:[12176993](#), PubMed:[14500380](#), PubMed:[27754931](#)). Can also act on sphingosylphosphorylcholine producing sphingosine-1-phosphate, a modulator of cell motility (PubMed:[14500380](#)). Can hydrolyze, in vitro, bis-pNPP, to some extent pNP-TMP, and barely ATP (PubMed:[12176993](#), PubMed:[15769751](#)). Involved in several motility-related processes such as angiogenesis and neurite outgrowth. Acts as an angiogenic factor by stimulating migration of smooth muscle cells and microtubule formation (PubMed:[11559573](#)). Stimulates migration of melanoma cells, probably via a pertussis toxin- sensitive G protein (PubMed:[1733949](#)). May have a role in induction of parturition (PubMed:[12176993](#)). Possible involvement in cell proliferation and adipose tissue development (Probable). Required for LPA production in activated platelets, cleaves the sn-1 lysophospholipids to generate sn-1 lysophosphatidic acids containing predominantly 18:2 and 20:4 fatty acids (PubMed:[21393252](#)). Shows a preference for the sn-1 to the sn-2 isomer of 1-O-alkyl-sn-glycero-3-phosphocholine (lyso-PAF) (PubMed:[21393252](#)).

#### Cellular Location

Secreted

#### Tissue Location

Detected in blood plasma (at protein level) (PubMed:[12176993](#), PubMed:[26371182](#)). Predominantly expressed in brain, placenta, ovary, and small intestine. Expressed in a number of carcinomas such as hepatocellular and prostate carcinoma, neuroblastoma and non-small-cell lung cancer. Expressed in body fluids such as plasma, cerebral spinal fluid (CSF), saliva, follicular and amniotic fluids. Not detected in leukocytes. Isoform 1 is more highly expressed in peripheral tissues than in the central nervous system (CNS) Adipocytes only express isoform 1. Isoform 3 is more highly expressed in the brain than in peripheral tissues.

## Background

---

ENPP2 functions as both a phosphodiesterase, which cleaves phosphodiester bonds at the 5' end of oligonucleotides, and a phospholipase, which catalyzes production of lysophosphatidic acid (LPA) in extracellular fluids. LPA evokes growth factor-like responses including stimulation of cell proliferation and chemotaxis. This protein stimulates the motility of tumor cells and has angiogenic properties, and its expression is upregulated in several kinds of carcinomas. The protein is secreted and further processed to make the biologically active form.

## References


---

Kawagoe H., Soma O., Goji J., Nishimura N., Narita M., Genomics 30:380-384(1995)  
 Nam S.W., Clair T., Kim Y.S., McMarlin A., Cancer Res. 61:6938-6944(2001)  
 The MGC Project Team Genome Res. 14:2121-2127(2004)

## Images

---

Anti-ENPP2 Antibody (Center) at 1:2000 dilution + Human brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 99 kDa  
 Blocking/Dilution buffer: 5% NFDm/TBST.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.