

APRT Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP2893b

Product Information

Application	IHC-P, WB, E
Primary Accession	P07741
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	19608
Antigen Region	143-170

Additional Information

Gene ID	353
Other Names	Adenine phosphoribosyltransferase, APRT, APRT
Target/Specificity	This APRT antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 143-170 amino acids from the C-terminal region of human APRT.
Dilution	IHC-P~~1:100~500 WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	APRT Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	APRT (HGNC:626)
Function	Catalyzes a salvage reaction resulting in the formation of AMP, that is energetically less costly than de novo synthesis.
Cellular Location	Cytoplasm.

Background


A conserved feature of APRT is the distribution of CpG dinucleotides. This enzyme catalyzes the formation of AMP and inorganic pyrophosphate from adenine and 5-phosphoribosyl-1-pyrophosphate (PRPP). It also produces adenine as a by-product of the polyamine biosynthesis pathway. A homozygous deficiency in this enzyme causes 2,8-dihydroxyadenine urolithiasis.

References


Silva,C.H., et. al.,J. Biomol. Struct. Dyn. 25 (6), 589-597 (2008)

Di Pietro,V., et. al., Clin. Biochem. 40 (1-2), 73-80 (2007)


Images



All lanes : Anti-APRT Antibody (C-term) at 1:1000 dilution
Lane 1: K562 whole cell lysate Lane 2: MCF-7 whole cell lysate
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 20 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



APRT Antibody (C-term) (Cat. #AP2893b) western blot analysis in MCF-7 cell line and rat kidney tissue lysates (35ug/lane). This demonstrates the APRT antibody detected the APRT protein (arrow).



Formalin-fixed and paraffin-embedded human hepatocarcinoma reacted with APRT Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

Citations

- [Adenine phosphoribosyltransferase \(APRT\) deficiency: identification of a novel nonsense mutation.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.