

# Phospho-H3(S28) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP3007a

## Product Information

---

Application	WB, E
Primary Accession	<a href="#">P68431</a>
Other Accession	<a href="#">Q71DJ3</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB5201
Calculated MW	15404

## Additional Information

---

Gene ID	8350;8351;8352;8353;8354;8355;8356;8357;8358;8968
Other Names	Histone H31, Histone H3/a, Histone H3/b, Histone H3/c, Histone H3/d, Histone H3/f, Histone H3/h, Histone H3/i, Histone H3/j, Histone H3/k, Histone H3/l, HIST1H3A, H3FA
Target/Specificity	This H3 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S28 of human H3.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Phospho-H3(S28) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	H3C1 ( <a href="#">HGNC:4766</a> )
Synonyms	H3FA, HIST1H3A
Function	Core component of nucleosome. Nucleosomes wrap and compact DNA into

chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

#### Cellular Location

Nucleus. Chromosome.

## Background

---

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3.


## References

---

- Lusic, M., et al., EMBO J. 22(24):6550-6561 (2003).  
Deng, L., et al., Virology 289(2):312-326 (2001).  
Deng, L., et al., Virology 277(2):278-295 (2000).  
El Kharroubi, A., et al., Mol. Cell. Biol. 18(5):2535-2544 (1998).  
Albig, W., et al., Hum. Genet. 101(3):284-294 (1997).

## Images

---



Western blot analysis of anti-Phospho-H3-pS28 Pab (Cat. #AP3007a) in HepG2 cell line lysate (35ug/lane). Mouse Phospho-H3-pS28(arrow) was detected using the purified Pab.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.