

Phospho-Bid(S65) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP3041a

Product Information

Application	WB, IHC-P, E
Primary Accession	P55957
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB7105
Calculated MW	21995

Additional Information

Gene ID	637
Other Names	BH3-interacting domain death agonist, p22 BID, BID, BH3-interacting domain death agonist p15, p15 BID, BH3-interacting domain death agonist p13, p13 BID, BH3-interacting domain death agonist p11, p11 BID, BID
Target/Specificity	This Bid Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S65 of human Bid.
Dilution	WB~1:1000 IHC-P~1:100~500 E~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Phospho-Bid(S65) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	BID
Function	Induces caspases and apoptosis (PubMed: 14583606). Counters the protective effect of BCL2 (By similarity).
Cellular Location	Cytoplasm. Mitochondrion membrane. Mitochondrion outer membrane.

Note=When uncleaved, it is predominantly cytoplasmic. [BH3-interacting domain death agonist p13]: Mitochondrion membrane {ECO:0000250|UniProtKB:P70444}. Note=Associated with the mitochondrial membrane. {ECO:0000250|UniProtKB:P70444} [Isoform 3]: Cytoplasm

Tissue Location

[Isoform 2]: Expressed in spleen, pancreas and placenta (at protein level).
[Isoform 4]: Expressed in lung and pancreas (at protein level).


Background

Bid is a death agonist that heterodimerizes with either agonist BAX or antagonist BCL2. The encoded protein is a member of the BCL-2 family of cell death regulators. Bid induces ICE-like proteases and apoptosis. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the major proteolytic product p15 Bid translocates to mitochondria where it triggers cytochrome c release.


References

- Liu, J., et al., *Biochem. Biophys. Res. Commun.* 330(3):865-870 (2005).
Broaddus, V.C., et al., *J. Biol. Chem.* 280(13):12486-12493 (2005).
Weng, C., et al., *J. Biol. Chem.* 280(11):10491-10500 (2005).
Gong, X.M., et al., *J. Biol. Chem.* 279(28):28954-28960 (2004).
Garcia-Saez, A.J., et al., *Biochemistry* 43(34):10930-10943 (2004).

Images




The anti-Phospho-Bid-S65 Pab (Cat. #AP3041a) is used in Western blot to detect Phospho-Bid-S65 in Jurkat tissue lysate



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma;



HC = hepatocarcinoma.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.