10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Phospho-CDC25A(S278) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP3047a

## Product Information

| | |
|---|---|
| **Application** | WB, IHC-P, E |
| **Primary Accession** | P30304 |
| **Reactivity** | Human |
| **Predicted** | Rabbit |
| **Host** | Rabbit |
| **Clonality** | Polyclonal |
| **Isotype** | Rabbit IgG |
| **Clone Names** | RB07909 |
| **Calculated MW** | 59087 |

## Additional Information

| | |
|---|---|
| **Gene ID** | 993 |
| **Other Names** | M-phase inducer phosphatase 1, Dual specificity phosphatase Cdc25A, CDC25A |
| **Target/Specificity** | This CDC25A Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S278 of human CDC25A. |
| **Dilution** | WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration. |
| **Format** | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| **Storage** | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| **Precautions** | Phospho-CDC25A(S278) Antibody is for research use only and not for use in diagnostic or therapeutic procedures. |

## Protein Information

| | |
|---|---|
| **Name** | CDC25A |
| **Function** | Tyrosine protein phosphatase which functions as a dosage- dependent inducer of mitotic progression (PubMed:12676925, PubMed:14559997, PubMed:1836978, PubMed:20360007). Directly dephosphorylates CDK1 and stimulates its kinase activity (PubMed:20360007). Also dephosphorylates |

CDK2 in complex with cyclin-E, in vitro (PubMed:20360007).


## Background

CDC25A is a member of the CDC25 family of phosphatases. CDC25A is required for progression from G1 to the S phase of the cell cycle. It activates the cyclin-dependent kinase CDC2 by removing two phosphate groups. CDC25A is specifically degraded in response to DNA damage, which prevents cells with chromosomal abnormalities from progressing through cell division. CDC25A is an oncogene, although its exact role in oncogenesis has not been demonstrated. Two transcript variants encoding different isoforms have been found for this gene.


## References


Ito, Y., et al., Int. J. Mol. Med. 13(3):431-435 (2004).
Nemoto, K., et al., Prostate 58(1):95-102 (2004).
Goloudina, A., et al., Cell Cycle 2(5):473-478 (2003).
Chen, M.S., et al., Mol. Cell. Biol. 23(21):7488-7497 (2003).
Chow, J.P., et al., Mol. Biol. Cell 14(10):3989-4002 (2003).


## Images




The anti-Phospho-CDC25A-S278 Pab (Cat. #AP3047a) is used in Western blot to detect Phospho-CDC25A-S278 in mouse heart tissue lysate




Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.


Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.