

Phospho-cJun(S63) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP3072a

Product Information

IHC-P, E <u>P05627, 077627</u> Mouse, Rat nal gG

Additional Information

Gene ID	3725
Other Names	Transcription factor AP-1, Activator protein 1, AP1, Proto-oncogene c-Jun, V-jun avian sarcoma virus 17 oncogene homolog, p39, JUN
Target/Specificity	This cJun Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S63 of human cJun.
Dilution	DB~~1:500 WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Phospho-cJun(S63) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	JUN
Function	Transcription factor that recognizes and binds to the AP-1 consensus motif 5'-TGA[GC]TCA-3' (PubMed: <u>10995748</u> , PubMed: <u>22083952</u>). Heterodimerizes

	with proteins of the FOS family to form an AP-1 transcription complex, thereby enhancing its DNA binding activity to the AP-1 consensus sequence 5'-TGA[GC]TCA-3' and enhancing its transcriptional activity (By similarity). Together with FOSB, plays a role in activation-induced cell death of T cells by binding to the AP-1 promoter site of FASLG/CD95L, and inducing its transcription in response to activation of the TCR/CD3 signaling pathway (PubMed:12618758). Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation (PubMed:17210646). Involved in activated KRAS-mediated transcriptional activation of USP28 in colorectal cancer (CRC) cells (PubMed:24623306).
Cellular Location	Nucleus.
Tissue Location	Expressed in the developing and adult prostate and prostate cancer cells.


Background

This gene for the cJun protein is the putative transforming gene of avian sarcoma virus 17. The protein is highly similar to the viral protein, and interacts directly with specific target DNA sequences to regulate gene expression. The gene for this protein is intronless and is mapped to 1p32-p31, a chromosomal region involved in both translocations and deletions in human malignancies.

References


Fang, D., et al., Proc. Natl. Acad. Sci. U.S.A. 101(41):14782-14787 (2004). Wang, Y., et al., Biochem. Biophys. Res. Commun. 323(1):9-16 (2004). Wehkamp, J., et al., Infect. Immun. 72(10):5750-5758 (2004). Gensch, E., et al., J. Biol. Chem. 279(37):39085-39093 (2004). Fujioka, S., et al., Mol. Cell. Biol. 24(17):7806-7819 (2004).

Images



Dot blot analysis of anti-Phospho-cJun-S63 Antibody (Cat.#AP3072a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

Western blot analysis of anti-Phospho-cJun-pS63 Pab (Cat. #AP3072a) in mouse kidney tissue lysate (35ug/lane). cJun-pS63 (arrow) was detected using the purified Pab.



Formali tissue re peroxid followed of this a relevand HC = he

Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.