10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Phospho-cJun(S63) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP3073a

Product Information

Application	WB, DB, IHC-P, E
Primary Accession	P05412
Reactivity	Human
Predicted	Mouse, Rat, Bovine, Canine, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB5193
Calculated MW	35676

Additional Information

Gene ID	3725
Other Names	Transcription factor AP-1, Activator protein 1, AP1, Proto-oncogene c-Jun, V-jun avian sarcoma virus 17 oncogene homolog, p39, JUN
Target/Specificity	This cJun Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S63 of human cJun.
Dilution	WB~~1:1000 DB~~1:500 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Phospho-cJun(S63) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	JUN
Function	Transcription factor that recognizes and binds to the AP-1 consensus motif 5'-TGA[GC]TCA-3' (PubMed:10995748, PubMed:22083952). Heterodimerizes with proteins of the FOS family to form an AP-1 transcription complex,

	thereby enhancing its DNA binding activity to the AP-1 consensus sequence 5'-TGA[GC]TCA-3' and enhancing its transcriptional activity (By similarity). Together with FOSB, plays a role in activation-induced cell death of T cells by binding to the AP-1 promoter site of FASLG/CD95L, and inducing its transcription in response to activation of the TCR/CD3 signaling pathway (PubMed:12618758). Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation (PubMed:17210646). Involved in activated KRAS-mediated transcriptional activation of USP28 in colorectal cancer (CRC) cells (PubMed:24623306). Binds to the USP28 promoter in colorectal cancer (CRC) cells (PubMed:24623306).
Cellular Location	Nucleus.
Tissue Location	Expressed in the developing and adult prostate and prostate cancer cells.

Background

The gene for cJun is the putative transforming gene of avian sarcoma virus 17. The cJun protein is a transcription factor highly similar to the viral protein, and interacts directly with specific target DNA sequences to regulate gene expression. The gene maps to 1p32-p31, a chromosomal region involved in both translocations and deletions in human malignancies.

References

Cheng, J., et al., J. Biol. Chem. 280(15):14492-14498 (2005).
Quan, T., et al., J. Biol. Chem. 280(9):8079-8085 (2005).
Bladh, L.G., et al., Mol. Pharmacol. 67(3):815-826 (2005).
DeNardo, D.G., et al., Mol. Endocrinol. 19(2):362-378 (2005).
Cheung, E., et al., Proc. Natl. Acad. Sci. U.S.A. 102(3):559-564 (2005).

Citations

- The DNMT1-associated lincRNA DACOR1 reprograms genome-wide DNA methylation in colon cancer.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.