

# Phospho-Wee1(S123) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP3284a

## Product Information

Application	WB, DB, IHC-P, E
Primary Accession	<a href="#">P30291</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	71597

## Additional Information

Gene ID	7465
Other Names	Wee1-like protein kinase, WEE1hu, Wee1A kinase, WEE1
Target/Specificity	This Wee1 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S123 of human Wee1.
Dilution	WB~~1:1000 DB~~1:500 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Phospho-Wee1(S123) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

Name	WEE1 {ECO:0000303 PubMed:8348613, ECO:0000312 HGNC:HGNC:12761}
Function	Acts as a negative regulator of entry into mitosis (G2 to M transition) by protecting the nucleus from cytoplasmically activated cyclin B1-complexed CDK1 before the onset of mitosis by mediating phosphorylation of CDK1 on 'Tyr-15' (PubMed: <a href="#">15070733</a> , PubMed: <a href="#">7743995</a> , PubMed: <a href="#">8348613</a> , PubMed: <a href="#">8428596</a> ). Specifically phosphorylates and inactivates cyclin B1-complexed CDK1 reaching a maximum during G2 phase and a minimum as cells enter M phase (PubMed: <a href="#">7743995</a> , PubMed: <a href="#">8348613</a> ,

PubMed:[8428596](#)). Phosphorylation of cyclin B1-CDK1 occurs exclusively on 'Tyr-15' and phosphorylation of monomeric CDK1 does not occur (PubMed:[7743995](#), PubMed:[8348613](#), PubMed:[8428596](#)). Its activity increases during S and G2 phases and decreases at M phase when it is hyperphosphorylated (PubMed:[7743995](#)). A correlated decrease in protein level occurs at M/G1 phase, probably due to its degradation (PubMed:[7743995](#)).

#### Cellular Location

Nucleus.

## Background

---

This gene encodes a nuclear protein, which is a tyrosine kinase belonging to the Ser/Thr family of protein kinases. This protein catalyzes the inhibitory tyrosine phosphorylation of CDC2/cyclin B kinase, and appears to coordinate the transition between DNA replication and mitosis by protecting the nucleus from cytoplasmically activated CDC2 kinase.


## References

---

Dai, X., et al., J. Invest. Dermatol. 122(6):1356-1364 (2004).  
Watanabe, N., et al., Proc. Natl. Acad. Sci. U.S.A. 101(13):4419-4424 (2004).  
Yoshida, T., et al., Ann. Oncol. 15(2):252-256 (2004).  
Kawasaki, H., et al., Oncogene 22(44):6839-6844 (2003).  
Yuan, H., et al., J. Virol. 77(3):2063-2070 (2003).

## Images

---



Dot blot analysis of Phospho-Wee1(S123) Antibody  
Phospho-specific Pab (Cat. AP3284a) on nitrocellulose  
membrane. 50ng of Phospho-peptide or Non  
Phospho-peptide per dot were adsorbed. Antibodies  
working concentration was 0.5ug per ml

## Citations

---

- [Persistence of the cell-cycle checkpoint kinase Wee1 in SadA- and SadB-deficient neurons disrupts neuronal polarity.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.