10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Phospho-ATF2(T339) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP3372a

Product Information

Application	WB, DB, E
Primary Accession	P15336
Reactivity	Human
Predicted	Mouse, Rat, Bovine, Canine, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB10901
Calculated MW	54537

Additional Information

Gene ID	1386
Other Names	Cyclic AMP-dependent transcription factor ATF-2, cAMP-dependent transcription factor ATF-2, Activating transcription factor 2, Cyclic AMP-responsive element-binding protein 2, CREB-2, cAMP-responsive element-binding protein 2, HB16, Histone acetyltransferase ATF2, cAMP response element-binding protein CRE-BP1, ATF2, CREB2, CREBP1
Target/Specificity	This ATF2 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S322 of human ATF2.
Dilution	WB~~1:1000 DB~~1:500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Phospho-ATF2(T339) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ATF2
Synonyms	CREB2, CREBP1

Function	Transcriptional activator which regulates the transcription of various genes, including those involved in anti-apoptosis, cell growth, and DNA damage response (PubMed:11932306, PubMed:15105425, PubMed:15916964, PubMed:22304920, PubMed:2529117). Dependent on its binding partner, binds to CRE (cAMP response element) consensus sequences (5'-TGACGTCA-3') or to AP-1 (activator protein 1) consensus sequences (5'-TGACTCA-3') (PubMed:11932306, PubMed:2529117). In the nucleus, contributes to global transcription and the DNA damage response, in addition to specific transcriptional activities that are related to cell development, proliferation and death (PubMed:22304920). In liver, acts as a regulator of glucose metabolism downstream of OR5V1 signaling, which promotes ATF2 activation, leading to expression of carnitine palmitoyltransferase CPT1A (By similarity). In the cytoplasm, interacts with and perturbs HK1- and VDAC1-containing complexes at the mitochondrial outer membrane, thereby impairing mitochondrial membrane potential, inducing mitochondrial leakage and promoting cell death (PubMed:22304920). The phosphorylated form (mediated by ATM) plays a role in the DNA damage response and is involved in the ionizing radiation (IR)-induced S phase checkpoint control and in the recruitment of the MRN complex into the IR-induced foci (IRIF) (PubMed:15916964). In concert with CUL3 and RBX1, promotes the degradation of KAT5 thereby attenuating its ability to acetylate and activate ATM (PubMed:18397884). Can elicit oncogenic or tumor suppressor activities depending on the tissue or cell type (PubMed:18677098).
Cellular Location	Nucleus. Cytoplasm Mitochondrion outer membrane Note=Shuttles between the cytoplasm and the nucleus and heterodimerization with JUN is essential for the nuclear localization (PubMed:22275354). Localization to the cytoplasm is observed under conditions of cellular stress and in disease states (PubMed:22304920) Localizes at the mitochondrial outer membrane in response to genotoxic stress (PubMed:22304920). Phosphorylation at Thr-52 is required for its nuclear localization and negatively regulates its mitochondrial localization (PubMed:22304920). Co-localizes with the MRN complex in the IR-induced foci (IRIF) (PubMed:15916964)
Tissue Location	Ubiquitously expressed, with more abundant expression in the brain.

Background


ATF2 is a transcription factor that is a member of the leucine zipper family of DNA binding proteins. This protein binds to the cAMP-responsive element (CRE), an octameric palindrome. The protein forms a homodimer or heterodimer with c-Jun and stimulates CRE-dependent transcription. The protein is also a histone acetyltransferase (HAT) that specifically acetylates histones H2B and H4 in vitro; thus it may represent a class of sequence-specific factors that activate transcription by direct effects on chromatin components.

References


Kravets, A., et al., J. Biol. Chem. 279(19):19916-19923 (2004).
Hong, S., et al., J. Biol. Chem. 279(17):16996-17003 (2004).
Averous, J., et al., J. Biol. Chem. 279(7):5288-5297 (2004).
Berger, A.J., et al., Cancer Res. 63(23):8103-8107 (2003).
Wen-Sheng, W., Oncogene 22(7):955-963 (2003).

Images

Dot blot analysis of Phospho-ATF2(S322)Antibody



Phospho-specific Pab (Cat. AP3372a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antobodies working concentration was 0. 5ug per ml.



Dot blot analysis of Phospho-ATF2-S322 Antibody (Cat. #AP3372a) and ATF2 Non Phospho-specific Pab on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.