

# Phospho-DNA-PK(T2609) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP3484a

## Product Information

---

Application	DB, E
Primary Accession	<a href="#">P78527</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB8113
Calculated MW	469089

## Additional Information

---

Gene ID	5591
Other Names	DNA-dependent protein kinase catalytic subunit, DNA-PK catalytic subunit, DNA-PKcs, DNPK1, p460, PRKDC, HYRC, HYRC1
Target/Specificity	This DNA-PK Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding T2609 of human DNA-PK.
Dilution	DB~~1:500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Phospho-DNA-PK(T2609) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	PRKDC
Synonyms	HYRC, HYRC1
Function	Serine/threonine-protein kinase that acts as a molecular sensor for DNA damage (PubMed: <a href="#">11955432</a> , PubMed: <a href="#">12649176</a> , PubMed: <a href="#">14734805</a> , PubMed: <a href="#">33854234</a> ). Involved in DNA non-homologous end joining (NHEJ)

required for double-strand break (DSB) repair and V(D)J recombination (PubMed:[11955432](#), PubMed:[12649176](#), PubMed:[14734805](#), PubMed:[33854234](#), PubMed:[34352203](#)). Must be bound to DNA to express its catalytic properties (PubMed:[11955432](#)). Promotes processing of hairpin DNA structures in V(D)J recombination by activation of the hairpin endonuclease artemis (DCLRE1C) (PubMed:[11955432](#)). Recruited by XRCC5 and XRCC6 to DNA ends and is required to (1) protect and align broken ends of DNA, thereby preventing their degradation, (2) and sequester the DSB for repair by NHEJ (PubMed:[11955432](#), PubMed:[12649176](#), PubMed:[14734805](#), PubMed:[15574326](#), PubMed:[33854234](#)). Acts as a scaffold protein to aid the localization of DNA repair proteins to the site of damage (PubMed:[11955432](#), PubMed:[12649176](#), PubMed:[14734805](#), PubMed:[15574326](#)). The assembly of the DNA-PK complex at DNA ends is also required for the NHEJ ligation step (PubMed:[11955432](#), PubMed:[12649176](#), PubMed:[14734805](#), PubMed:[15574326](#)). Found at the ends of chromosomes, suggesting a further role in the maintenance of telomeric stability and the prevention of chromosomal end fusion (By similarity). Also involved in modulation of transcription (PubMed:[11955432](#), PubMed:[12649176](#), PubMed:[14734805](#), PubMed:[15574326](#)). As part of the DNA-PK complex, involved in the early steps of ribosome assembly by promoting the processing of precursor rRNA into mature 18S rRNA in the small-subunit processome (PubMed:[32103174](#)). Binding to U3 small nucleolar RNA, recruits PRKDC and XRCC5/Ku86 to the small-subunit processome (PubMed:[32103174](#)). Recognizes the substrate consensus sequence [ST]-Q (PubMed:[11955432](#), PubMed:[12649176](#), PubMed:[14734805](#), PubMed:[15574326](#)). Phosphorylates 'Ser-139' of histone variant H2AX, thereby regulating DNA damage response mechanism (PubMed:[14627815](#), PubMed:[16046194](#)). Phosphorylates ASF1A, DCLRE1C, c-Abl/ABL1, histone H1, HSPCA, c-jun/JUN, p53/TP53, PARP1, POU2F1, DHX9, FH, SRF, NHEJ1/XLF, XRCC1, XRCC4, XRCC5, XRCC6, WRN, MYC and RFA2 (PubMed:[10026262](#), PubMed:[10467406](#), PubMed:[11889123](#), PubMed:[12509254](#), PubMed:[14599745](#), PubMed:[14612514](#), PubMed:[14704337](#), PubMed:[15177042](#), PubMed:[1597196](#), PubMed:[16397295](#), PubMed:[18644470](#), PubMed:[2247066](#), PubMed:[2507541](#), PubMed:[26237645](#), PubMed:[26666690](#), PubMed:[28712728](#), PubMed:[29478807](#), PubMed:[30247612](#), PubMed:[8407951](#), PubMed:[8464713](#), PubMed:[9139719](#), PubMed:[9362500](#)). Can phosphorylate C1D not only in the presence of linear DNA but also in the presence of supercoiled DNA (PubMed:[9679063](#)). Ability to phosphorylate p53/TP53 in the presence of supercoiled DNA is dependent on C1D (PubMed:[9363941](#)). Acts as a regulator of the phosphatidylinositol 3-kinase/protein kinase B signal transduction by mediating phosphorylation of 'Ser-473' of protein kinase B (PKB/AKT1, PKB/AKT2, PKB/AKT3), promoting their activation (PubMed:[15262962](#)). Contributes to the determination of the circadian period length by antagonizing phosphorylation of CRY1 'Ser-588' and increasing CRY1 protein stability, most likely through an indirect mechanism (By similarity). Plays a role in the regulation of DNA virus-mediated innate immune response by assembling into the HDP-RNP complex, a complex that serves as a platform for IRF3 phosphorylation and subsequent innate immune response activation through the cGAS-STING pathway (PubMed:[28712728](#)). Also regulates the cGAS-STING pathway by catalyzing phosphorylation of CGAS, thereby impairing CGAS oligomerization and activation (PubMed:[33273464](#)). Also regulates the cGAS-STING pathway by mediating phosphorylation of PARP1 (PubMed:[35460603](#)).

## Cellular Location

Nucleus. Nucleus, nucleolus. Cytoplasm, cytosol

## Background

Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein

kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains. The STE group (homologs of yeast Sterile 7, 11, 20 kinases) consists of 50 kinases related to the mitogen-activated protein kinase (MAPK) cascade families (Ste7/MAP2K, Ste11/MAP3K, and Ste20/MAP4K). MAP kinase cascades, consisting of a MAPK and one or more upstream regulatory kinases (MAPKKs) have been best characterized in the yeast pheromone response pathway. Pheromones bind to Stem cell surface receptors and activate yeast MAPK pathway.


## References

---

- Goudelock, D.M., et al., J. Biol. Chem. 278(32):29940-29947 (2003).  
Ding, Q., et al., Mol. Cell. Biol. 23(16):5836-5848 (2003).  
Lucero, H., et al., J. Biol. Chem. 278(24):22136-22143 (2003).  
Calsou, P., et al., J. Mol. Biol. 326(1):93-103 (2003).  
Karpova, A.Y., et al., Proc. Natl. Acad. Sci. U.S.A. 99(5):2818-2823 (2002).

## Images

---



Dot blot analysis of anti-DNA-PK-pT2609 Pab (RB08113) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.