10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Phospho-GFAP(S8) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP3562a

Product Information

Application	DB, E
Primary Accession	P14136
Other Accession	NP_002046
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB15197
Calculated MW	49880

Additional Information

Gene ID	2670
Other Names	Glial fibrillary acidic protein, GFAP, GFAP
Target/Specificity	This GFAP Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S8 of human GFAP.
Dilution	DB~~1:500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Phospho-GFAP(S8) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	GFAP
Function	GFAP, a class-III intermediate filament, is a cell-specific marker that, during the development of the central nervous system, distinguishes astrocytes from other glial cells.
Cellular Location	Cytoplasm. Note=Associated with intermediate filaments

Tissue Location	Expressed in cells lacking fibronectin.


Background

GFAP is one of the major intermediate filament proteins of mature astrocytes. It is used as a marker to distinguish astrocytes from other glial cells during development. Mutations in this gene cause Alexander disease, a rare disorder of astrocytes in the central nervous system.

References

Quintanar, J.L., et al., Parasitol. Res. 90(4):261-263 (2003).
Shiroma, N., et al., Brain Dev. 25(2):116-121 (2003).
Nielsen, A.L., et al., J. Biol. Chem. 277(33):29983-29991 (2002).
Namekawa, M., et al., Ann. Neurol. 52(6):779-785 (2002).
Lopez-Egido, J., et al., Exp. Cell Res. 278(2):175-183 (2002).

Images



Dot blot analysis of anti-Phospho-GFAP-S8 Antibody (Cat.#AP3562a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

Citations

- Analysis of Chaperone-Mediated Autophagy.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.