

# Phospho-GFAP(S8) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP3562a

## Product Information

---

<b>Application</b>	DB, E
<b>Primary Accession</b>	<a href="#">P14136</a>
<b>Other Accession</b>	<a href="#">NP_002046</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB15197
<b>Calculated MW</b>	49880

## Additional Information

---

<b>Gene ID</b>	2670
<b>Other Names</b>	Glial fibrillary acidic protein, GFAP, GFAP
<b>Target/Specificity</b>	This GFAP Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S8 of human GFAP.
<b>Dilution</b>	DB~~1:500 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	Phospho-GFAP(S8) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	GFAP
<b>Function</b>	GFAP, a class-III intermediate filament, is a cell-specific marker that, during the development of the central nervous system, distinguishes astrocytes from other glial cells.
<b>Cellular Location</b>	Cytoplasm. Note=Associated with intermediate filaments

## Tissue Location

Expressed in cells lacking fibronectin.

## Background

---

GFAP is one of the major intermediate filament proteins of mature astrocytes. It is used as a marker to distinguish astrocytes from other glial cells during development. Mutations in this gene cause Alexander disease, a rare disorder of astrocytes in the central nervous system.


## References

---

Quintanar, J.L., et al., Parasitol. Res. 90(4):261-263 (2003).  
Shiroma, N., et al., Brain Dev. 25(2):116-121 (2003).  
Nielsen, A.L., et al., J. Biol. Chem. 277(33):29983-29991 (2002).  
Namekawa, M., et al., Ann. Neurol. 52(6):779-785 (2002).  
Lopez-Egido, J., et al., Exp. Cell Res. 278(2):175-183 (2002).

## Images

---



Dot blot analysis of anti-Phospho-GFAP-S8 Antibody (Cat.#AP3562a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

## Citations

---

- [Analysis of Chaperone-Mediated Autophagy.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.