

# Phospho-ANTXR1(Y425) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP3603a

## **Product Information**

**Application** DB, E **Primary Accession** Q9H6X2

Other Accession OOPMD2, O9CZ52

Reactivity Human
Predicted Mouse, Rat
Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Clone Names RB16635
Calculated MW 62789

## **Additional Information**

**Gene ID** 84168

Other Names Anthrax toxin receptor 1, Tumor endothelial marker 8, ANTXR1, ATR, TEM8

**Target/Specificity** This ANTXR1 Antibody is generated from rabbits immunized with a KLH

conjugated synthetic phosphopeptide corresponding to amino acid residues

surrounding Y425 of human ANTXR1.

**Dilution** DB~~1:500 E~~Use at an assay dependent concentration.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** Phospho-ANTXR1(Y425) Antibody is for research use only and not for use in

diagnostic or therapeutic procedures.

#### **Protein Information**

Name ANTXR1 {ECO:0000303 | PubMed:22912819,

ECO:0000312 | HGNC:HGNC:21014}

**Function** Plays a role in cell attachment and migration. Interacts with extracellular

matrix proteins and with the actin cytoskeleton and thereby plays an important role in normal extracellular matrix (ECM) homeostasis. Mediates

adhesion of cells to type 1 collagen and gelatin, reorganization of the actin cytoskeleton and promotes cell spreading. Plays a role in the angiogenic response of cultured umbilical vein endothelial cells. May also act as a receptor for PLAU. Upon ligand binding, stimulates the phosphorylation of EGFR and ERK1/2 (PubMed:30241478).

#### **Cellular Location**

Cell membrane; Single-pass type I membrane protein. Cell projection, lamellipodium membrane; Single-pass type I membrane protein. Cell projection, filopodium membrane; Single-pass type I membrane protein. Note=At the membrane of lamellipodia and at the tip of actin-enriched filopodia (PubMed:16762926). Colocalizes with actin at the base of lamellipodia (PubMed:16762926)

**Tissue Location** 

Detected in umbilical vein endothelial cells (at protein level). Highly expressed in tumor endothelial cells


# **Background**

ANTXR1 is a type I transmembrane protein and is a tumor-specific endothelial marker that has been implicated in colorectal cancer. This protein has been shown to also be a docking protein or receptor for Bacillus anthracis toxin, the causative agent of the disease, anthrax. The binding of the protective antigen (PA) component, of the tripartite anthrax toxin, to this receptor protein mediates delivery of toxin components to the cytosol of cells. Once inside the cell, the other two components of anthrax toxin, edema factor (EF) and lethal factor (LF) disrupt normal cellular processes.

## References

Werner, E., J. Biol. Chem. 281 (32), 23227-23236 (2006) Abrami, L., J. Cell Biol. 172 (2), 309-320 (2006) Bradley, K.A., Nature 414 (6860), 225-229 (2001)

# **Images**



Dot blot analysis of anti-Phospho-ANTXR1-pY425 Antibody (Cat.#AP3603a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.