

# Phospho-CCNB2(S392) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP3839a

## Product Information

---

Application	DB, E
Primary Accession	<a href="#">O95067</a>
Other Accession	<a href="#">NP_004692.1</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB42149
Calculated MW	45282

## Additional Information

---

Gene ID	9133
Other Names	G2/mitotic-specific cyclin-B2, CCNB2
Target/Specificity	This CCNB2 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S392 of human CCNB2.
Dilution	DB~~1:500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Phospho-CCNB2(S392) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	CCNB2
Function	Essential for the control of the cell cycle at the G2/M (mitosis) transition.

## Background

---

Cyclin B2 is a member of the cyclin family, specifically the B-type cyclins. The B-type cyclins, B1 and B2, associate with p34cdc2 and are essential components of the cell cycle regulatory machinery. B1 and B2 differ in their subcellular localization. Cyclin B1 co-localizes with microtubules, whereas cyclin B2 is primarily associated with the Golgi region. Cyclin B2 also binds to transforming growth factor beta RII and thus cyclin B2/cdc2 may play a key role in transforming growth factor beta-mediated cell cycle control.


## References

---

Cunningham, J.M., et al. Br. J. Cancer 101(8):1461-1468(2009)  
Haraguchi, T., et al. Fertil. Steril. 91 (4 SUPPL), 1424-1426 (2009) :  
De Martino, I., et al. Cancer Res. 69(5):1844-1850(2009)  
Bellanger, S., et al. Oncogene 26(51):7175-7184(2007)  
Stav, D., et al. Int. J. Biol. Markers 22(2):108-113(2007)

## Images

---



Dot blot analysis of CCNB2 Antibody (Phospho S392)  
Phospho-specific Pab (Cat. #AP3839a) on nitrocellulose  
membrane. 50ng of Phospho-peptide or Non  
Phospho-peptide per dot were adsorbed. Antibody  
working concentrations are 0.6ug per ml.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.