

# Phospho-CCNB1(S35) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP3882a

## Product Information

---

<b>Application</b>	DB, WB, E
<b>Primary Accession</b>	<a href="#">P14635</a>
<b>Other Accession</b>	<a href="#">NP_114172.1</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB42141
<b>Calculated MW</b>	48337

## Additional Information

---

<b>Gene ID</b>	891
<b>Other Names</b>	G2/mitotic-specific cyclin-B1, CCNB1, CCNB
<b>Target/Specificity</b>	This CCNB1 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S35 of human CCNB1.
<b>Dilution</b>	DB~~1:500 WB~~1:1100 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	Phospho-CCNB1(S35) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	CCNB1
<b>Synonyms</b>	CCNB
<b>Function</b>	Essential for the control of the cell cycle at the G2/M (mitosis) transition.
<b>Cellular Location</b>	Cytoplasm. Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center,


## Background

The protein encoded by this gene is a regulatory protein involved in mitosis. The gene product complexes with p34(cdc2) to form the maturation-promoting factor (MPF). Two alternative transcripts have been found, a constitutively expressed transcript and a cell cycle-regulated transcript, that is expressed predominantly during G2/M phase. The different transcripts result from the use of alternate transcription initiation sites. [provided by RefSeq].


## References

Kreis, N.N., et al. *Oncogene* 29(41):5591-5603(2010)  
 van Zon, W., et al. *J. Cell Biol.* 190(4):587-602(2010)  
 Harley, M.E., et al. *EMBO J.* 29(14):2407-2420(2010)  
 Olson, J.E., et al. *Breast Cancer Res. Treat.* (2010) In press :  
 Nantajit, D., et al. *PLoS ONE* 5 (8), E12341 (2010) :

## Images



Western blot analysis of lysate from HeLa cells(from left to right),untreated or treated with Nocodazole at 50ng/ml,using Phospho-CCNB1-S35 Antibody (Cat. #AP3882a) or CCNB1-S9 Antibody.Lysate at 15ug per lane.AP3882a was diluted at 1:1100 dilution at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody.



Dot blot analysis of CCNB1 Antibody (Phospho S35) Phospho-specific Pab (Cat. #AP3882a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.6ug per ml.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.