

PAQR3 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP4717b

Product Information

Application	WB, E
Primary Accession	Q6TCH7
Other Accession	Q6TCG8
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB24961
Calculated MW	36217
Antigen Region	242-270

Additional Information

Gene ID	152559
Other Names	Progestin and adipoQ receptor family member 3, Progestin and adipoQ receptor family member III, Raf kinase trapping to Golgi, RKTG, PAQR3
Target/Specificity	This PAQR3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 242-270 amino acids from the C-terminal region of human PAQR3.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PAQR3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PAQR3 (HGNC:30130)
Function	Golgi-scaffold protein which modulates its interactors activities by anchoring them to the Golgi apparatus (PubMed: 18547165 , PubMed: 26311497).

Functions as a spatial regulator of RAF1 kinase by sequestering it to the Golgi apparatus (PubMed:[18547165](#)). Acts as a positive regulator of cholesterol biosynthesis by mediating the anchoring of the SCAP:SREBP complex in the Golgi apparatus, thereby promoting SCAP:SREBF2 complex formation, potentiating SREBF2 and SREBF1 processing and enhancing lipid synthesis (PubMed:[26311497](#)). Also regulates PPARA and PPARG functions by mediating their interaction with E3 ubiquitin ligases, such as STUB1 or HUWE1, leading to their polyubiquitination and proteasome-mediated degradation (By similarity).

Cellular Location

Golgi apparatus membrane; Multi-pass membrane protein

Tissue Location

Widely expressed in a range of tissues.

Background

PAQR3 is function as a spatial regulator of RAF1 kinase by sequestering it to the Golgi.


References

Jiang, Y., et al. Mol. Cell. Biol. 30(1):78-90(2010)

Garitaonandia, I., et al. J. Recept. Signal Transduct. Res. 29(1):67-73(2009)

Luo, X., et al. Biochem. J. 414(3):399-406(2008)

Images



All lanes: Anti-PAQR3 Antibody (C-term) at 1:500 dilution + Mouse testis lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 34 KDa Blocking/Dilution buffer: 5% NFDM/TBST.

Citations

- [Expression of adipokines in osteoarthritis osteophytes and their effect on osteoblasts.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.