

ARHGAP25 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP4865c

Product Information

Application IHC-P, WB, E
Primary Accession P42331
Other Accession Q8BYW1
Reactivity Human, Mouse

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Clone Names RB22842
Calculated MW 73435
Antigen Region 193-219

Additional Information

Gene ID 9938

Other Names Rho GTPase-activating protein 25, Rho-type GTPase-activating protein 25,

ARHGAP25, KIAA0053

Target/SpecificityThis ARHGAP25 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 193-219 amino acids from the Central

region of human ARHGAP25.

Dilution IHC-P~~1:100~500 WB~~1:1000 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions ARHGAP25 Antibody (Center) is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name ARHGAP25

Synonyms KIAA0053

Function GTPase activator for the Rho-type GTPases by converting them to an


Background

ARHGAP25 encode negative regulators of Rho GTPases (see ARHA; MIM 165390), which are implicated in actin remodeling, cell polarity, and cell migration.


References

Takefuji, M., et al. J. Hum. Genet. 55(1):42-49(2010) Marroni, F., et al. Circ Cardiovasc Genet 2(4):322-328(2009) Katoh, M., et al. Int. J. Mol. Med. 14(2):333-338(2004)

Images



Anti-ARHGAP25 Antibody (Center) at 1:1000 dilution + Jurkat whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 73 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



ARHGAP25 Antibody (Center) (Cat. #AP4865c) immunohistochemistry analysis in formalin fixed and paraffin embedded human tonsil tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the ARHGAP25 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.