

# HIF1A Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP4879d

## Product Information

---

|                   |                        |
|-------------------|------------------------|
| Application       | WB, FC, IHC-P, E       |
| Primary Accession | <a href="#">Q16665</a> |
| Reactivity        | Human                  |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |
| Isotype           | Rabbit IgG             |
| Clone Names       | RB23012                |
| Calculated MW     | 92670                  |

## Additional Information

---

|                    |                                                                                                                                                                                                                                                                         |
|--------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Gene ID            | 3091                                                                                                                                                                                                                                                                    |
| Other Names        | Hypoxia-inducible factor 1-alpha, HIF-1-alpha, HIF1-alpha, ARNT-interacting protein, Basic-helix-loop-helix-PAS protein MOP1, Class E basic helix-loop-helix protein 78, bHLHe78, Member of PAS protein 1, PAS domain-containing protein 8, HIF1A, BHLHE78, MOP1, PASD8 |
| Target/Specificity | This HIF1A antibody is generated from rabbits immunized with HIF1A recombinant protein.                                                                                                                                                                                 |
| Dilution           | WB~~1:1000 FC~~1:10~50 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.                                                                                                                                                                                     |
| Format             | Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.                                                                                             |
| Storage            | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.                                                                                                                                 |
| Precautions        | HIF1A Antibody is for research use only and not for use in diagnostic or therapeutic procedures.                                                                                                                                                                        |

## Protein Information

---

|          |                                                                                                                                                                                                                                                                                                        |
|----------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Name     | HIF1A {ECO:0000303 PubMed:7539918, ECO:0000312 HGNC:HGNC:4910}                                                                                                                                                                                                                                         |
| Function | Functions as a master transcriptional regulator of the adaptive response to hypoxia (PubMed: <a href="#">11292861</a> , PubMed: <a href="#">11566883</a> , PubMed: <a href="#">15465032</a> , PubMed: <a href="#">16973622</a> , PubMed: <a href="#">17610843</a> , PubMed: <a href="#">18658046</a> , |

PubMed:[20624928](#), PubMed:[22009797](#), PubMed:[30125331](#), PubMed:[9887100](#)). Under hypoxic conditions, activates the transcription of over 40 genes, including erythropoietin, glucose transporters, glycolytic enzymes, vascular endothelial growth factor, HILPDA, and other genes whose protein products increase oxygen delivery or facilitate metabolic adaptation to hypoxia (PubMed:[11292861](#), PubMed:[11566883](#), PubMed:[15465032](#), PubMed:[16973622](#), PubMed:[17610843](#), PubMed:[20624928](#), PubMed:[22009797](#), PubMed:[30125331](#), PubMed:[9887100](#)). Plays an essential role in embryonic vascularization, tumor angiogenesis and pathophysiology of ischemic disease (PubMed:[22009797](#)). Heterodimerizes with ARNT; heterodimer binds to core DNA sequence 5'-TACGTG-3' within the hypoxia response element (HRE) of target gene promoters (By similarity). Activation requires recruitment of transcriptional coactivators such as CREBBP and EP300 (PubMed:[16543236](#), PubMed:[9887100](#)). Activity is enhanced by interaction with NCOA1 and/or NCOA2 (PubMed:[10594042](#)). Interaction with redox regulatory protein APEX1 seems to activate CTAD and potentiates activation by NCOA1 and CREBBP (PubMed:[10202154](#), PubMed:[10594042](#)). Involved in the axonal distribution and transport of mitochondria in neurons during hypoxia (PubMed:[19528298](#)).

#### Cellular Location

Cytoplasm. Nucleus. Nucleus speckle {ECO:0000250|UniProtKB:Q61221}. Note=Colocalizes with HIF3A in the nucleus and speckles (By similarity). Cytoplasmic in normoxia, nuclear translocation in response to hypoxia (PubMed:[9822602](#)) {ECO:0000250|UniProtKB:Q61221, ECO:0000269|PubMed:[9822602](#)}

#### Tissue Location

Expressed in most tissues with highest levels in kidney and heart. Overexpressed in the majority of common human cancers and their metastases, due to the presence of intratumoral hypoxia and as a result of mutations in genes encoding oncoproteins and tumor suppressors. A higher level expression seen in pituitary tumors as compared to the pituitary gland.

## Background


Hypoxia-inducible factor-1 (HIF1) is a transcription factor found in mammalian cells cultured under reduced oxygen tension that plays an essential role in cellular and systemic homeostatic responses to hypoxia. HIF1 is a heterodimer composed of an alpha subunit and a beta subunit. The beta subunit has been identified as the aryl hydrocarbon receptor nuclear translocator (ARNT). This gene encodes the alpha subunit of HIF-1. Overexpression of a natural antisense transcript (aHIF) of this gene has been shown to be associated with nonpapillary renal carcinomas.

## References



Lee, M.N., et al. J. Natl. Cancer Inst. 102(6):426-442(2010)  
 Mayer, A., et al. Adv. Exp. Med. Biol. 662, 399-405 (2010)  
 Brouwer, E., et al. Clin. Exp. Rheumatol. 27(6):945-951(2009)

## Images


All lanes: Anti-HIF1A Antibody at 1:1000 dilution Lane 1: NIH/3T3 whole cell lysate Lane 2: Raji whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 93 KDa Blocking/Dilution buffer: 5% NFDm/TBST.



Western blot analysis of HIF1A Antibody (Cat. #AP4879d) in K562 cell line lysates (35ug/lane). HIF1A (arrow) was detected using the purified Pab.



HIF1A Antibody (Cat. #AP4879d) IHC analysis in formalin fixed and paraffin embedded lung carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the HIF1A Antibody for immunohistochemistry. Clinical relevance has not been evaluated.



HIF1A Antibody (Cat. #AP4879d) flow cytometric analysis of K562 cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.