

CC50B Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP4968a

Product Information

Application	WB, IHC-P, FC, E
Primary Accession	Q3MIR4
Other Accession	NP_001017970
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB25806
Calculated MW	38941
Antigen Region	42-71

Additional Information

Gene ID	161291
Other Names	Cell cycle control protein 50B, P4-ATPase flippase complex beta subunit TMEM30B, Transmembrane protein 30B, TMEM30B, CDC50B
Target/Specificity	This CC50B antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 42-71 amino acids from the N-terminal region of human CC50B.
Dilution	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	CC50B Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TMEM30B
Synonyms	CDC50B

Function

Accessory component of a P4-ATPase flippase complex which catalyzes the hydrolysis of ATP coupled to the transport of aminophospholipids from the outer to the inner leaflet of various membranes and ensures the maintenance of asymmetric distribution of phospholipids. Phospholipid translocation also seems to be implicated in vesicle formation and in uptake of lipid signaling molecules. The beta subunit may assist in binding of the phospholipid substrate (Probable). Can mediate the export of alpha subunits ATP8A1, ATP8B1, ATP8B2 and ATP8B4 from the ER to the plasma membrane.


Cellular Location

Cell membrane; Multi-pass membrane protein


References

Katoh, Y., et al. Oncol. Rep. 12(4):939-943(2004)


Images



Western blot analysis of CC50B Antibody (N-term) (Cat. #AP4968a) in MDA-MB435 cell line lysates (35ug/lane). CC50B (arrow) was detected using the purified Pab.



CC50B Antibody (N-term) (Cat. #AP4968a) IHC analysis in formalin fixed and paraffin embedded human breast carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the CC50B Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



CC50B Antibody (N-term) (Cat. #AP4968a) flow cytometric analysis of MDA-MB435 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.