

# Cyclin D2 Antibody (Ab-280)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP50014

## Product Information

---

Application	WB
Primary Accession	<a href="#">P30279</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	polyclonal
Calculated MW	33067

## Additional Information

---

Gene ID	894
Other Names	G1/S-specific cyclin-D2, CCND2
Dilution	WB~~ 1:500-1:1000
Format	Rabbit IgG in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.
Storage Conditions	-20°C

## Protein Information

---

Name	CCND2 {ECO:0000303   PubMed:1386336, ECO:0000312   HGNC:HGNC:1583}
Function	Regulatory component of the cyclin D2-CDK4 (DC) complex that phosphorylates and inhibits members of the retinoblastoma (RB) protein family including RB1 and regulates the cell-cycle during G(1)/S transition (PubMed: <a href="#">18827403</a> , PubMed: <a href="#">8114739</a> ). Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complex and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase (PubMed: <a href="#">18827403</a> , PubMed: <a href="#">8114739</a> ). Hypophosphorylates RB1 in early G(1) phase (PubMed: <a href="#">18827403</a> , PubMed: <a href="#">8114739</a> ). Cyclin D-CDK4 complexes are major integrators of various mitogenic and antimitogenic signals (PubMed: <a href="#">18827403</a> , PubMed: <a href="#">8114739</a> ).
Cellular Location	Nucleus. Cytoplasm. Nucleus membrane. Note=Cyclin D-CDK4 complexes accumulate at the nuclear membrane and are then translocated into the nucleus through interaction with KIP/CIP family members

## Background

---

Regulatory component of the cyclin D2-CDK4 (DC) complex that phosphorylates and inhibits members of the retinoblastoma (RB) protein family including RB1 and regulates the cell-cycle during G(1)/S transition. Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complex and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase. Hypophosphorylates RB1 in early G(1) phase. Cyclin D-CDK4 complexes are major integrators of various mitogenic and antimitogenic signals. Also substrate for SMAD3, phosphorylating SMAD3 in a cell-cycle-dependent manner and repressing its transcriptional activity. Component of the ternary complex, cyclin D2/CDK4/CDKN1B, required for nuclear translocation and activity of the cyclin D-CDK4 complex (By similarity).


## References

---

Xiong Y.,et al.Genomics 13:575-584(1992).  
Palmero I.,et al.Oncogene 8:1049-1054(1993).  
Miyajima N.,et al.Submitted (MAR-1993) to the EMBL/GenBank/DDBJ databases.  
Kalnine N.,et al.Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.  
Ota T.,et al.Nat. Genet. 36:40-45(2004).

## Images

---



Western blot analysis of lysate from RD cell line,using Cyclin D2 Antibody (Ab-280)(B8336). B8336 was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody.Lysate at 35ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.