

Cytochrome P450 2A6 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP50056

Product Information

Application WB
Primary Accession P11509
Reactivity Human
Host Rabbit
Clonality polyclonal
Calculated MW 56517

Additional Information

Gene ID 1548

Other Names Cytochrome P450 2A6, 11413-, 4-cineole 2-exo-monooxygenase, CYPIIA6,

Coumarin 7-hydroxylase, Cytochrome P450 IIA3, Cytochrome P450(I), CYP2A6,

CYP2A3

Dilution WB~~ 1:500-1:1000

Format Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4,

150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.

Storage Conditions -20°C

Protein Information

Name CYP2A6

Synonyms CYP2A3

Function Exhibits a high coumarin 7-hydroxylase activity. Can act in the hydroxylation

of the anti-cancer drugs cyclophosphamide and ifosphamide. Competent in the metabolic activation of aflatoxin B1. Constitutes the major nicotine C-oxidase. Acts as a 1,4-cineole 2-exo- monooxygenase. Possesses low

phenacetin O-deethylation activity.

Cellular Location Endoplasmic reticulum membrane; Peripheral membrane protein. Microsome

membrane; Peripheral membrane protein

Tissue Location Liver.


Background

Exhibits a high coumarin 7-hydroxylase activity. Can act in the hydroxylation of the anti-cancer drugs cyclophosphamide and ifosphamide. Competent in the metabolic activation of aflatoxin B1. Constitutes the major nicotine C-oxidase. Acts as a 1,4- cineole 2-exo-monooxygenase. Possesses low phenacetin O-deethylation activity.

References

Miles J.S., et al. Nucleic Acids Res. 17:2907-2917(1989). Yamano S., et al. Nucleic Acids Res. 17:4888-4888(1989). Yamano S., et al. Biochemistry 29:1322-1329(1990). Zhuge J., et al. Submitted (SEP-1999) to the EMBL/GenBank/DDBJ databases. Ota T., et al. Nat. Genet. 36:40-45(2004).

Images



Western blot analysis of lysates from H.liver cell line ,using Cytochrome P450 2A6 Antibody(C12258). C12258 was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody.Lysates at 35ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.