

RAD18 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP50198

Product Information

Application	WB
Primary Accession	Q9NS91
Reactivity	Human, Rat
Host	Rabbit
Clonality	polyclonal
Calculated MW	56223

Additional Information

Gene ID	56852
Other Names	E3 ubiquitin-protein ligase RAD18, 632-, Postreplication repair protein RAD18, hHR18, hRAD18, RING finger protein 73, RAD18, RNF73
Dilution	WB~~ 1:1000
Format	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.
Storage Conditions	-20°C

Protein Information

Name	RAD18
Synonyms	RNF73
Function	E3 ubiquitin-protein ligase involved in postreplication repair of UV-damaged DNA. Postreplication repair functions in gap- filling of a daughter strand on replication of damaged DNA. Associates to the E2 ubiquitin conjugating enzyme UBE2B to form the UBE2B-RAD18 ubiquitin ligase complex involved in mono-ubiquitination of DNA- associated PCNA on 'Lys-164'. Has ssDNA binding activity.
Cellular Location	Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=Associates with chromatin (PubMed:25931565). Colocalizes with SLF1 in the nucleus and to centrosomes (PubMed:15632077). Relocalizes with SLF1 to nuclear foci in response to DNA damage (PubMed:22036607). Accumulates with the SLF1- SLF2 and SMC5-SMC6 complexes at replication-coupled DNA interstrand repair and DNA double-strand breaks (DSBs) sites on chromatin in a ubiquitin-dependent manner (PubMed:25931565)


Background

E3 ubiquitin-protein ligase involved in postreplication repair of UV-damaged DNA. Postreplication repair functions in gap- filling of a daughter strand on replication of damaged DNA. Associates to the E2 ubiquitin conjugating enzyme UBE2B to form the UBE2B-RAD18 ubiquitin ligase complex involved in mono-ubiquitination of DNA-associated PCNA on 'Lys-164'. Has ssDNA binding activity.

References

Tateishi S.,et al.Proc. Natl. Acad. Sci. U.S.A. 97:7927-7932(2000).
Xin H.,et al.Nucleic Acids Res. 28:2847-2854(2000).
Jang Y.,et al.Submitted (JUL-2000) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Muzny D.M.,et al.Nature 440:1194-1198(2006).

Images



Western blot analysis of lysates from HeLa, MCF-7 cell line (from left to right), using RAD18 Antibody (C15564). C15564 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35 µg per lane.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.