

# TNF12 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP50579

## Product Information

Application	WB, IF, IHC
Primary Accession	<a href="#">O43508</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Calculated MW	27216

## Additional Information

Gene ID	8742
Other Names	Tumor necrosis factor ligand superfamily member 12, APO3 ligand, TNF-related weak inducer of apoptosis, TWEAK, Tumor necrosis factor ligand superfamily member 12, membrane form, Tumor necrosis factor ligand superfamily member 12, secreted form, TNFSF12, APO3L, DR3LG
Dilution	WB~~ 1:1000 IF~~1:100 IHC~~1:50-1:100
Format	Rabbit IgG in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.
Storage Conditions	-20°C

## Protein Information

Name	TNFSF12
Synonyms	APO3L, DR3LG
Function	Binds to FN14 and possibly also to TNFSF12/APO3. Weak inducer of apoptosis in some cell types. Mediates NF-kappa-B activation. Promotes angiogenesis and the proliferation of endothelial cells. Also involved in induction of inflammatory cytokines. Promotes IL8 secretion.
Cellular Location	Cell membrane; Single-pass type II membrane protein [Isoform TWE-PRIL]; Cell membrane; Single-pass membrane protein
Tissue Location	Highly expressed in adult heart, pancreas, skeletal muscle, brain, colon, small intestine, lung, ovary, prostate, spleen, lymph node, appendix and peripheral blood lymphocytes. Low expression in kidney, testis, liver, placenta, thymus and bone marrow. Also detected in fetal kidney, liver, lung and brain


## Background

Binds to FN14 and possibly also to TNFRSF12/APO3. Weak inducer of apoptosis in some cell types. Mediates NF-kappa-B activation. Promotes angiogenesis and the proliferation of endothelial cells. Also involved in induction of inflammatory cytokines. Promotes IL8 secretion.


## References

Chicheportiche Y.,et al.J. Biol. Chem. 272:32401-32410(1997).  
Marsters S.A.,et al.Curr. Biol. 8:525-528(1998).  
Pradet-Balade B.,et al.EMBO J. 21:5711-5720(2002).  
Clark H.F.,et al.Genome Res. 13:2265-2270(2003).  
Zody M.C.,et al.Nature 440:1045-1049(2006).


## Images



Western blot analysis of lysates from 293, HUVEC cell line (from left to right), using TNF12 Antibody (AP50579). AP50579 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35 µg per lane.



Immunofluorescence analysis of HeLa cells, using TNF12 antibody.



Immunohistochemistry analysis of paraffin-embedded human brain tissue using TNF12 antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.