

G3BP2 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP50602

Product Information

Application	WB
Primary Accession	Q9UN86
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	polyclonal
Calculated MW	54121

Additional Information

Gene ID	9908
Other Names	Ras GTPase-activating protein-binding protein 2, G3BP-2, GAP SH3 domain-binding protein 2, G3BP2, KIAA0660
Dilution	WB~~ 1:500
Format	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.
Storage Conditions	-20°C

Protein Information

Name	G3BP2 {ECO:0000303 PubMed:23279204, ECO:0000312 HGNC:HGNC:30291}
Function	Scaffold protein that plays an essential role in cytoplasmic stress granule formation which acts as a platform for antiviral signaling (PubMed: 23279204 , PubMed: 32302570 , PubMed: 32302571 , PubMed: 32302572). Plays an essential role in stress granule formation (PubMed: 27022092 , PubMed: 32302570 , PubMed: 32302571 , PubMed: 32302572 , PubMed: 35977029). Stress granules are membraneless compartments that store mRNAs and proteins, such as stalled translation pre-initiation complexes, in response to stress (PubMed: 32302570 , PubMed: 32302571 , PubMed: 32302572). Promotes formation of stress granules phase-separated membraneless compartment by undergoing liquid-liquid phase separation (LLPS) upon unfolded RNA-binding: functions as a molecular switch that triggers RNA-dependent LLPS in response to a rise in intracellular free RNA concentrations (By similarity).
Cellular Location	Cytoplasm. Cytoplasm, Stress granule


Background

Probable scaffold protein that may be involved in mRNA transport (Potential).

References

Kennedy D.,et al.Submitted (APR-1999) to the EMBL/GenBank/DDBJ databases.
Ishikawa K.,et al.DNA Res. 5:169-176(1998).
Guitard E.,et al.Submitted (FEB-1998) to the EMBL/GenBank/DDBJ databases.
Kennedy D.,et al.Submitted (MAR-1998) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).

Images



Western blot analysis of lysates from HeLa, 293 cell line (from left to right), using G3BP2 Antibody (AP50602). AP50602 was diluted at 1:500 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.

Image not found : 201408/AP50602_2.jpg

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.