

NDUFV2 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP50682

Product Information

Application	WB
Primary Accession	P19404
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	polyclonal
Calculated MW	27392

Additional Information

Gene ID	4729
Other Names	NADH dehydrogenase [ubiquinone] flavoprotein 2, mitochondrial, NADH-ubiquinone oxidoreductase 24 kDa subunit, NDUFV2
Dilution	WB~~1:1000
Format	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.
Storage Conditions	-20°C

Protein Information

Name	NDUFV2 (HGNC:7717)
Function	Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which catalyzes electron transfer from NADH through the respiratory chain, using ubiquinone as an electron acceptor (Probable). Parts of the peripheral arm of the enzyme, where the electrons from NADH are accepted by flavin mononucleotide (FMN) and then passed along a chain of iron-sulfur clusters by electron tunnelling to the final acceptor ubiquinone (Probable). Contains one iron-sulfur cluster (Probable).
Cellular Location	Mitochondrion inner membrane {ECO:0000250 UniProtKB:P04394}; Peripheral membrane protein {ECO:0000250 UniProtKB:P04394}; Matrix side {ECO:0000250 UniProtKB:P04394}

Background

Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) that is

believed to belong to the minimal assembly required for catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone (By similarity).

References


Pilkington S.J.,et al.Biochemistry 28:3257-3264(1989).

Murray J.,et al.J. Biol. Chem. 278:13619-13622(2003).

Burkard T.R.,et al.BMC Syst. Biol. 5:17-17(2011).

Ogura M.,et al.Biochem. J. 447:281-289(2012).

Images



Western blot analysis of lysate from HeLa cell line,using NDUFV2 Antibody(AP50682). AP50682 was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody.Lysate at 35ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.