

NARG1 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP50688

Product Information

Application	WB
Primary Accession	Q9BXJ9
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Calculated MW	101272

Additional Information

Gene ID	80155
Other Names	N-alpha-acetyltransferase 15, NatA auxiliary subunit, Gastric cancer antigen Ga19, N-terminal acetyltransferase, NMDA receptor-regulated protein 1, Protein tubedown-1, Tbdn100, NAA15, GA19, NARG1, NATH, TBDN100
Dilution	WB~~ 1:1000
Format	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.
Storage Conditions	-20°C

Protein Information

Name	NAA15
Synonyms	GA19, NARG1, NATH, TBDN100
Function	Auxillary subunit of N-terminal acetyltransferase complexes which display alpha (N-terminal) acetyltransferase (NAT) activity (PubMed: 15496142 , PubMed: 20154145 , PubMed: 29754825 , PubMed: 32042062). The NAT activity may be important for vascular, hematopoietic and neuronal growth and development (PubMed: 15496142). Required to control retinal neovascularization in adult ocular endothelial cells (PubMed: 11687548). In complex with XRCC6 and XRCC5 (Ku80), up-regulates transcription from the osteocalcin promoter (PubMed: 12145306).
Cellular Location	Cytoplasm. Nucleus. Note=Mainly cytoplasmic, nuclear in some cases. Present in the free cytosolic and cytoskeleton- bound polysomes, but not in the membrane-bound polysomes
Tissue Location	Expressed at high levels in testis and in ocular endothelial cells. Also found in

brain (corpus callosum), heart, colon, bone marrow and at lower levels in most adult tissues, including thyroid, liver, pancreas, mammary and salivary glands, lung, ovary, urogenital system and upper gastrointestinal tract. Overexpressed in gastric cancer, in papillary thyroid carcinomas and in a Burkitt lymphoma cell line (Daudi). Specifically suppressed in abnormal proliferating blood vessels in eyes of patients with proliferative diabetic retinopathy.


Background

The NAA10-NAA15 complex displays alpha (N-terminal) acetyltransferase activity that may be important for vascular, hematopoietic and neuronal growth and development. Required to control retinal neovascularization in adult ocular endothelial cells. In complex with XRCC6 and XRCC5 (Ku80), up-regulates transcription from the osteocalcin promoter.

References

- Line A.,et al.Br. J. Cancer 86:1824-1830(2002).
Willis D.M.,et al.J. Biol. Chem. 277:37280-37291(2002).
Fluge O.,et al.Oncogene 21:5056-5068(2002).
He Y.G.,et al.Sheng Wu Hua Xue Yu Sheng Wu Wu Li Xue Bao 34:353-357(2002).
Ota T.,et al.Nat. Genet. 36:40-45(2004).

Images



Western blot analysis of lysate from 293T cell line, using NARG1 Antibody (AP50688). AP50688 was diluted at 1:1000. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.