

SNX4 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP50705

Product Information

Application	WB
Primary Accession	<u>095219</u>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	polyclonal
Calculated MW	51909

Additional Information

Gene ID	8723
Other Names	Sorting nexin-4, SNX4
Dilution	WB~~1:500
Format	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.
Storage Conditions	-20°C

Protein Information

Name	SNX4 {ECO:0000303 PubMed:9819414, ECO:0000312 HGNC:HGNC:11175}
Function	Involved in the regulation of endocytosis and in several stages of intracellular trafficking (PubMed: <u>12668730</u> , PubMed: <u>17994011</u> , PubMed: <u>32513819</u> , PubMed: <u>33468622</u>). Plays a role in recycling endocytosed transferrin receptor and prevent its degradation (PubMed: <u>17994011</u>). Involved in autophagosome assembly by regulating trafficking and recycling of phospholipid scramblase ATG9A (PubMed: <u>32513819</u> , PubMed: <u>33468622</u>).
Cellular Location	Early endosome membrane; Peripheral membrane protein; Cytoplasmic side. Note=Also detected on a juxtanuclear endocytic recycling compartment (ERC).

Background

May be involved in several stages of intracellular trafficking. Plays a role in recycling endocytosed transferrin receptor and prevent its degradation.

References

Haft C.R.,et al.Mol. Cell. Biol. 18:7278-7287(1998). Ota T.,et al.Nat. Genet. 36:40-45(2004). Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases. Traer C.J.,et al.Nat. Cell Biol. 9:1370-1380(2007). Dephoure N.,et al.Proc. Natl. Acad. Sci. U.S.A. 105:10762-10767(2008).

Images



Western blot analysis of lysate from HepG2 cell line, using SNX4 Antibody(AP50705). AP50705 was diluted at 1:500. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35 ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.