10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-IFN omega Antibody

Catalog # AP50726

Product Information

Application	WB, IF/IC
Primary Accession	P05000
Reactivity	Human, Mouse, Rat, Mk
Host	Rabbit
Clonality	Polyclonal
Calculated MW	22319

Additional Information

Gene ID	3467
Other Names	Interferon omega-1; Interferon alpha-II-1
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human IFN omega. The exact sequence is proprietary.
Dilution	WB~~ 1:1000 IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	IFNW1
Cellular Location	Secreted.


References

Capon D.J.,et al.Mol. Cell. Biol. 5:768-779(1985).
Adolf G.R.,et al.Biochim. Biophys. Acta 1089:167-174(1991).
Humphray S.J.,et al.Nature 429:369-374(2004).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Hauptmann R.,et al.Nucleic Acids Res. 13:4739-4749(1985).

Images



Western blot analysis of IFN omega expression in HEK293T (A), NIHC3T3 (B), H9C2 (C) whole cell lysates. (Predicted band size: 22 kD; Observed band size: 22 kD)



Immunofluorescent analysis of IFN omega staining in H460 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.