

POU4F1 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP50748

Product Information

Application	WB
Primary Accession	Q01851
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	polyclonal
Calculated MW	42697

Additional Information

Gene ID	5457
Other Names	POU domain, class 4, transcription factor 1, Brain-specific homeobox/POU domain protein 3A, Brain-3A, Brn-3A, Homeobox/POU domain protein RDC-1, Oct-T1, POU4F1, BRN3A, RDC1
Dilution	WB~~1:1000
Format	Rabbit IgG in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.
Storage Conditions	-20°C

Protein Information

Name	POU4F1 (HGNC:9218)
Function	Multifunctional transcription factor with different regions mediating its different effects. Acts by binding (via its C-terminal domain) to sequences related to the consensus octamer motif 5'- ATGCAAAT-3' in the regulatory regions of its target genes. Regulates the expression of specific genes involved in differentiation and survival within a subset of neuronal lineages. It has been shown that activation of some of these genes requires its N-terminal domain, maybe through a neuronal-specific cofactor. Activates BCL2 expression and protects neuronal cells from apoptosis (via the N-terminal domain). Induces neuronal process outgrowth and the coordinate expression of genes encoding synaptic proteins. Exerts its major developmental effects in somatosensory neurons and in brainstem nuclei involved in motor control. Stimulates the binding affinity of the nuclear estrogen receptor ESR1 to DNA estrogen response element (ERE), and hence modulates ESR1-induced transcriptional activity. May positively regulate POU4F2 and POU4F3. Regulates dorsal root ganglion sensory neuron specification and axonal projection into the spinal cord. Plays a role in TNFSF11-mediated terminal

osteoclast differentiation. Negatively regulates its own expression interacting directly with a highly conserved autoregulatory domain surrounding the transcription initiation site.

Cellular Location

Nucleus. Cytoplasm {ECO:0000250 | UniProtKB:P17208}

Tissue Location

Expressed in the brain and the retina. Present in the developing brain, spinal cord and eye.


Background

Probable transcription factor which may play a role in the regulation of specific gene expression within a subset of neuronal lineages. May play a role in determining or maintaining the identities of a small subset of visual system neurons.

References

Bhargava A.K.,et al.Proc. Natl. Acad. Sci. U.S.A. 90:10260-10264(1993).
Xiang M.,et al.J. Neurosci. 15:4762-4785(1995).
Dunham A.,et al.Nature 428:522-528(2004).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Collum R.G.,et al.Nucleic Acids Res. 20:4919-4925(1992).

Images



Western blot analysis of lysate from SW480 cell line,using POU4F1 Antibody(AP50748). AP50748 was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody.Lysate at 35ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.