

TAIP-2 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP50827

Product Information

Application	WB
Primary Accession	Q8WYN3
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Calculated MW	64900

Additional Information

Gene ID	80034
Other Names	Cysteine/serine-rich nuclear protein 3, CSRNP-3, Protein FAM130A2, TGF-beta-induced apoptosis protein 2, TAIP-2, CSRNP3, FAM130A2, TAIP2
Dilution	WB~~ 1:1000
Format	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.
Storage Conditions	-20°C

Protein Information

Name	CSRNP3
Synonyms	FAM130A2, TAIP2
Function	Binds to the consensus sequence 5'-AGAGTG-3' and has transcriptional activator activity. Plays a role in apoptosis (By similarity).
Cellular Location	Nucleus.


Background

Binds to the consensus sequence 5'-AGAGTG-3' and has transcriptional activator activity. Plays a role in apoptosis (By similarity).

References

Akiyama N.,et al.Submitted (JUN-2001) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Hillier L.W.,et al.Nature 434:724-731(2005).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Sjoeblom T.,et al.Science 314:268-274(2006).

Images



Western blot analysis of lysate from PC-3 cell line,using TAIP-2 Antibody,was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody.Lysate at 35ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.