

SLC24A4 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP50838

Product Information

Application	WB
Primary Accession	Q8NFF2
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	polyclonal
Calculated MW	69042

Additional Information

Gene ID	123041
Other Names	Sodium/potassium/calcium exchanger 4, Na(+)/K(+)/Ca(2+)-exchange protein 4, Solute carrier family 24 member 4, SLC24A4, NCKX4
Dilution	WB~~ 1:1000
Format	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.
Storage Conditions	-20°C

Protein Information

Name	SLC24A4 (HGNC:10978)
Function	Calcium, potassium:sodium antiporter that transports 1 Ca(2+) and 1 K(+) in exchange for 4 Na(+) (PubMed: 12379639 , PubMed: 26631410). Controls the rapid response termination and proper regulation of adaptation in olfactory sensory neurons (OSNs) which subsequently influences how odor information is encoded and perceived (By similarity). May play a role in calcium transport during amelogenesis (PubMed: 23375655 , PubMed: 24621671).
Cellular Location	Cell membrane; Multi-pass membrane protein. Cytoplasm {ECO:0000250 UniProtKB:Q8CGQ8}
Tissue Location	Expressed abundantly in all regions of the brain, aorta, lung and thymus (PubMed:12379639). Expressed at lower levels in the stomach and intestine (PubMed:12379639)

Background

Transports 1 Ca(2+) and 1 K(+) in exchange for 4 Na(+). Controls the rapid response termination and proper regulation of adaptation in olfactory sensory neurons (OSNs) which subsequently influences how odor information is encoded and perceived. May play a role in calcium transport during amelogenesis (By similarity).

References

Li X.-F.,et al.J. Biol. Chem. 277:48410-48417(2002).


Ota T.,et al.Nat. Genet. 36:40-45(2004).

Heilig R.,et al.Nature 421:601-607(2003).

Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

Bechtel S.,et al.BMC Genomics 8:399-399(2007).

Images



Western blot analysis of lysate from human lung tissue lysate,using SLC24A4 Antibody, was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody.Lysate at 35ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.