

# **AQP4** Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP50937

## **Product Information**

Application WB
Primary Accession P55087
Reactivity Human, Rat
Host Rabbit
Clonality Polyclonal
Calculated MW 34830

## **Additional Information**

Gene ID 361

Other Names Aquaporin-4, AQP-4, Mercurial-insensitive water channel, MIWC, WCH4, AQP4

**Dilution** WB~~ 1:500

Format Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.09% (W/V)

sodium azide and 52% glycerol

**Storage** Store at -20 °C.Stable for 12 months from date of receipt

### **Protein Information**

Name AQP4

**Function** Forms a water-specific channel (PubMed: 19383790, PubMed: 7559426,

PubMed:8601457). Plays an important role in brain water homeostasis (PubMed:37143309). It is involved in glymphatic solute transport and is required for a normal rate of water exchange across the blood brain interface. Required for normal levels of cerebrospinal fluid influx into the brain cortex and parenchyma along paravascular spaces that surround penetrating arteries, and for normal drainage of interstitial fluid along paravenous drainage pathways. Thereby, it is required for normal clearance of solutes from the brain interstitial fluid, including soluble beta-amyloid peptides derived from APP. Plays a redundant role in urinary water

homeostasis and urinary concentrating ability (By similarity).

**Cellular Location** Cell membrane; Multi-pass membrane protein. Basolateral cell membrane

{ECO:0000250 | UniProtKB:P55088}; Multi-pass membrane protein. Endosome membrane {ECO:0000250 | UniProtKB:P47863}. Cell membrane, sarcolemma;

Multi-pass membrane protein. Cell projection

{ECO:0000250 | UniProtKB:P47863}. Note=Activation of the vasopressin receptor AVPR1A triggers AQP4 phosphorylation at Ser-180 and promotes its

internalization from the cell membrane. Detected on brain astrocyte processes and astrocyte endfeet close to capillaries {ECO:0000250|UniProtKB:P47863}

#### **Tissue Location**

Detected in skeletal muscle (PubMed:29055082). Detected in stomach, along the glandular base region of the fundic gland (at protein level) (PubMed:8601457). Detected in brain, lung and skeletal muscle, and at much lower levels in heart and ovary (PubMed:7559426, PubMed:8601457).


# **Background**

Forms a water-specific channel. Osmoreceptor which regulates body water balance and mediates water flow within the central nervous system.

## References

Yang B.,et al.J. Biol. Chem. 270:22907-22913(1995). Misaka T.,et al.FEBS Lett. 381:208-212(1996). Lu M.,et al.Proc. Natl. Acad. Sci. U.S.A. 93:10908-10912(1996). Nusbaum C.,et al.Nature 437:551-555(2005). Lanciotti A.,et al.Hum. Mol. Genet. 21:2166-2180(2012).

# **Images**



Western blot analysis of lysates from mouse RAW264.7,Hela,rat PC-12,U-87 MG cell line (from left to right), using AQP4 Antibody(AP50937). AP50937 was diluted at 1:500 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.Lysates at 20ug per lane.

# **Citations**

• Atorvastatin ameliorates early brain injury through inhibition of apoptosis and ER stress in a rat model of subarachnoid hemorrhage.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.