

ABCG1 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP50949

Product Information

Application	WB
Primary Accession	P45844
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	75592

Additional Information

Gene ID	9619
Other Names	ATP-binding cassette sub-family G member 1, ATP-binding cassette transporter 8, White protein homolog, ABCG1, ABC8, WHT1
Target/Specificity	KLH conjugated synthetic peptide derived from human ABCG1
Dilution	WB~~ 1:1000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	ABCG1 (HGNC:73)
Synonyms	ABC8, WHT1
Function	Catalyzes the efflux of phospholipids such as sphingomyelin, cholesterol and its oxygenated derivatives like 7beta- hydroxycholesterol and this transport is coupled to hydrolysis of ATP (PubMed: 17408620 , PubMed: 24576892). The lipid efflux is ALB-dependent (PubMed: 16702602). Is an active component of the macrophage lipid export complex. Could also be involved in intracellular lipid transport processes. The role in cellular lipid homeostasis may not be limited to macrophages. Prevents cell death by transporting cytotoxic 7beta-hydroxycholesterol (PubMed: 17408620).
Cellular Location	Endoplasmic reticulum membrane; Multi-pass membrane protein. Golgi apparatus membrane; Multi-pass membrane protein. Cell membrane Note=Predominantly localized in the intracellular compartments mainly associated with the endoplasmic reticulum (ER) and Golgi membranes

Tissue Location

Expressed in several tissues. Expressed in macrophages; expression is increased in macrophages from patients with Tangier disease.


Background

Transporter involved in macrophage lipid homeostasis. Is an active component of the macrophage lipid export complex. Could also be involved in intracellular lipid transport processes. The role in cellular lipid homeostasis may not be limited to macrophages.

References

Chen H.M.,et al.Am. J. Hum. Genet. 59:66-75(1996).
Berry A.,et al.Genomics 68:22-29(2000).
Porsch-Oezcueruemez M.,et al.J. Biol. Chem. 276:12427-12433(2001).
Lorkowski S.,et al.Biochem. Biophys. Res. Commun. 280:121-131(2001).
Kennedy M.A.,et al.J. Biol. Chem. 276:39438-39447(2001).

Images



Anti-ABCG1 Antibody at 1:1000 dilution + MCF-7 whole cell lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 76 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.

Citations

- [Murine Beta-Amyloid \(1-42\) Oligomers Disrupt Endothelial Barrier Integrity and VEGFR Signaling via Activating Astrocytes to Release Deleterious Soluble Factors](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.