

# Adenylate Cyclase 5/6 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP50965

#### **Product Information**

Application WB Primary Accession 095622

**Reactivity** Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW138908

#### **Additional Information**

Gene ID 111

**Other Names** Adenylate cyclase type 5, ATP pyrophosphate-lyase 5, Adenylate cyclase type

V, Adenylyl cyclase 5, ADCY5

**Target/Specificity** KLH-conjugated synthetic peptide encompassing a sequence within the

C-term region of human Adenylate Cyclase 5/6. The exact sequence is

proprietary.

**Dilution** WB~~ 1:1000

Format 0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

**Storage** Store at -20 °C.Stable for 12 months from date of receipt

#### **Protein Information**

Name ADCY5

**Function** Catalyzes the formation of the signaling molecule cAMP in response to

G-protein signaling (PubMed:<u>15385642</u>, PubMed:<u>24700542</u>, PubMed:<u>26206488</u>). Mediates signaling downstream of ADRB1 (PubMed:<u>24700542</u>). Regulates the increase of free cytosolic Ca(2+) in response to increased blood glucose levels and contributes to the regulation

of Ca(2+)-dependent insulin secretion (PubMed:24740569).

**Cell ular Location**Cell membrane; Multi-pass membrane protein. Cell projection, cilium

{ECO:0000250 | UniProtKB:P84309}

**Tissue Location** Detected in pancreas islets (at protein level). Expressed in the brain, with high

expression in the corpus striatum (PubMed:26085604).

## **Background**

This is a membrane-bound, calcium-inhibitable adenylyl cyclase.

### References

Ota T.,et al.Nat. Genet. 36:40-45(2004).


Muzny D.M.,et al.Nature 440:1194-1198(2006).

Ludwig M.G.,et al.J. Recept. Signal Transduct. 22:79-110(2002).

Raimundo S.,et al.Clin. Chim. Acta 285:155-161(1999).

Ding Q.,et al.Mol. Pharmacol. 66:921-928(2004).

## **Images**



Anti-Adenylate Cyclase 5/6 Antibodyat 1:1000 dilution + PC-12 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L),Peroxidase conjugated at 1/10000 dilution Predicted band size: 139 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.