10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Ataxin 2 Antibody

Catalog # AP51006

Product Information

Application	WB, IHC, IF/IC
Primary Accession	Q99700
Reactivity	Human, Mouse, Mk
Host	Rabbit
Clonality	Polyclonal
Calculated MW	140283

Additional Information

Gene ID	6311
Other Names	ATX2; SCA2; TNRC13; Ataxin-2; Spinocerebellar ataxia type 2 protein; Trinucleotide repeat-containing gene 13 protein
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Ataxin 2. The exact sequence is proprietary.
Dilution	WB~~ 1:2000 IHC~~1:100~500 IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	ATXN2
Synonyms	ATX2, SCA2, TNRC13
Function	Involved in EGFR trafficking, acting as negative regulator of endocytic EGFR internalization at the plasma membrane.
Cellular Location	Cytoplasm.
Tissue Location	Expressed in the brain, heart, liver, skeletal muscle, pancreas and placenta. Isoform 1 is predominant in the brain and spinal cord. Isoform 4 is more abundant in the cerebellum. In the brain, broadly expressed in the amygdala, caudate nucleus, corpus callosum, hippocampus, hypothalamus, substantia nigra, subthalamic nucleus and thalamus.


References

Pulst S.-M.,et al.Nat. Genet. 14:269-276(1996).
Sanpei K.,et al.Nat. Genet. 14:277-284(1996).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Scherer S.E.,et al.Nature 440:346-351(2006).
Imbert G.,et al.Nat. Genet. 14:285-291(1996).


Images



Western blot analysis of Ataxin 2 expression in HEK293T (A), PC3 (B), NIN3T3 (C) whole cell lysates. (Predicted band size: 140 kD; Observed band size: 170 kD)



Immunohistochemical analysis of Ataxin 2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of Ataxin 2 staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.