

Caspase 2 p18 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51040

Product Information

Application	WB
Primary Accession	P42575
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	50685

Additional Information

Gene ID	835
Other Names	Caspase-2, CASP-2, Neural precursor cell expressed developmentally down-regulated protein 2, NEDD-2, Protease ICH-1, Caspase-2 subunit p18, Caspase-2 subunit p13, Caspase-2 subunit p12, CASP2, ICH1, NEDD2
Target/Specificity	KLH conjugated synthetic peptide derived from human Caspase 2 p18
Dilution	WB~~ 1:1000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	CASP2
Synonyms	ICH1, NEDD2
Function	Is a regulator of the cascade of caspases responsible for apoptosis execution (PubMed: 11156409 , PubMed: 15073321 , PubMed: 8087842). Might function by either activating some proteins required for cell death or inactivating proteins necessary for cell survival (PubMed: 15073321). Associates with PIDD1 and CRADD to form the PIDDosome, a complex that activates CASP2 and triggers apoptosis in response to genotoxic stress (PubMed: 15073321).
Tissue Location	Expressed at higher levels in the embryonic lung, liver and kidney than in the heart and brain. In adults, higher level expression is seen in the placenta, lung, kidney, and pancreas than in the heart, brain, liver and skeletal muscle

Background

Involved in the activation cascade of caspases responsible for apoptosis execution. Might function by either activating some proteins required for cell death or inactivating proteins necessary for cell survival.

References

Wang L.,et al.Cell 78:739-750(1994).


Droin N.,et al.Cancer Res. 60:7039-7047(2000).

Halleck A.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Hillier L.W.,et al.Nature 424:157-164(2003).

Images



All lanes : Anti-Caspase 2 p18 Antibody at 1:1000 dilution
Lane 1: Jurkat whole cell lysates Lane 2: HeLa whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 51 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.